Risk factors for fragility fracture in middle age. A prospective population-based study of 33,000 men and women

**A. H. Holmberg · O. Johnell · J. Nilsson · K. Åkesson ·
P. M. Nilsson · G. Berglund**

Published online: 30 June 2006

© International Osteoporosis Foundation and National Osteoporosis Foundation 2006

The online version of the original article can be found at: <http://dx.doi.org/10.1007/s00198-006-0137-7>

A. H. Holmberg · O. Johnell · J. Nilsson · K. Åkesson
Clinical and Molecular Research Unit,
Department of Clinical Sciences Malmö, Lund University,
Malmö University Hospital,
SE-205 02 Malmö, Sweden

P. M. Nilsson · G. Berglund
Internal Medicine Research Unit,
Department of Clinical Sciences Malmö, Lund University,
Malmö University Hospital,
SE-205 02 Malmö, Sweden

K. Åkesson (✉)
Department of Orthopaedics, Malmö University Hospital,
20502 Malmö, Sweden
e-mail: kristina_akesson@med.lu.se

The wrong publication was listed as reference [24]. The correct reference [24] is as follows:

Reference

24. Jonsson B, Gardsell P, Johnell O, Redlund-Johnell I, Sembo I (1994) Remembering fractures: fracture registration and recall in southern Sweden. *J Epidemiol Community Health* 48(5):489–490