

Use of anti-depressants and the risk of fracture of the hip or femur

M. W. M. van den Brand · S. Pouwels · M. M. Samson · T. P. van Staa · B. Thio · C. Cooper · H. G. M. Leufkens · A. C. G. Egberts · H. J. J. Verhaar · F. de Vries

Published online: 27 March 2009

© International Osteoporosis Foundation and National Osteoporosis Foundation 2009

Erratum to: Osteoporos Int DOI 10.1007/s00198-009-0849-6

The names of the second and third authors were given in the wrong order. The correct order of authors is as given above.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00198-009-0849-6>.

M. W. M. van den Brand (✉) · M. M. Samson · H. J. J. Verhaar
Department of Geriatric Medicine,
University Medical Centre Utrecht,
Heidelberglaan 100,
3584 CX Utrecht, The Netherlands
e-mail: mwm.vandenbrand@wanadoo.nl

S. Pouwels · T. P. van Staa · B. Thio · H. G. M. Leufkens ·
A. C. G. Egberts · F. de Vries (✉)
Division of Pharmacoepidemiology and Pharmacotherapy,
Utrecht Institute for Pharmaceutical Sciences, Universiteit Utrecht,
Sorbonnelaan 16,
3584 CA Utrecht, The Netherlands
e-mail: f.devries@uu.nl

T. P. van Staa · C. Cooper · F. de Vries
Medical Research Council Epidemiology Resource Centre,
University of Southampton, Southampton General Hospital,
Southampton, UK

T. P. van Staa · F. de Vries
General Practice Research Database,
Medicines and Healthcare products Regulatory Agency,
19 Elmslane, SW8 5NQ,
London, UK

C. Cooper
Institute of Musculoskeletal Sciences, University of Oxford,
Oxford, UK

A. C. G. Egberts
Department of Clinical Pharmacy,
University Medical Centre Utrecht,
Heidelberglaan 100, 3584 CX,
Utrecht, The Netherlands