ERRATUM

Erratum to: A retrospective analysis of nonresponse to daily teriparatide treatment

R. Niimi¹ · T. Kono¹ · A. Nishihara¹ · M. Hasegawa² · T. Kono¹ · A. Sudo²

Published online: 17 May 2016

© International Osteoporosis Foundation and National Osteoporosis Foundation 2016

Erratum to: Osteoporos Int

DOI 10.1007/s00198-016-3581-z

The first sentence of the second paragraph in the section headed “Changes in the bone turnover markers’ concentrations in response to teriparatide treatment” should have read as follows: “In the nonresponders, the mean (SD) uNTX levels increased to 73 nmol BCE/mmol Cr (59) at 4 months, 87 nmol BCE/mmol Cr (64) at 8 months, 84 nmol BCE/mmol Cr (53) at 12 months, 86 nmol BCE/mmol Cr (64) at 16 months, 80 nmol BCE/mmol Cr (74) at 20 months, and 77 nmol BCE/mmol Cr (99) at 24 months (all $p < 0.01$ vs. baseline; Wilcoxon signed-rank test) (Fig. 2b)”.

In addition, both panels Fig. 1 contained errors. The amended figure is given here.

The authors apologise for any inconvenience caused.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00198-016-3581-z>.

✉ R. Niimi
furikakefuri@hotmail.co.jp

¹ Department of Orthopaedic Surgery, Tomidahama Hospital, 26-14, Tomidahamacho, Yokkaichi, Mie 510-8008, Japan

² Department of Orthopaedic Surgery, Graduate School of Medicine, Mie University, Tsu, Japan