Manori J. Silva · Dana B. Barr · John A. Reidy
Kayoko Kato · Nicole A. Malek · Carolyn C. Hodge
Donald Hurtz III · Antonia M. Calafat
Larry L. Needham · John W. Brock

Glucuronidation patterns of common urinary and serum monoester phthalate metabolites

Published online: 13 April 2005
© Springer-Verlag 2005

Arch Toxicol (2003) 77:561–567

Page 565, second column, second paragraph should read: The percentage of free mEP, mBP, mBzP and

mEHP found in urine remained scattered with geometric means 71, 6, 7 and 16% respectively, and showed no apparent increase with increased concentration of total phthalate as should occur if enzyme saturation occurred.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00204-003-0486-3>

M. J. Silva (✉) · D. B. Barr · J. A. Reidy · K. Kato
N. A. Malek · C. C. Hodge · A. M. Calafat · L. L. Needham
J. W. Brock
Division of Laboratory Sciences, National Center
for Environmental Health, Centers for Disease Control
and Prevention, 4770 Buford Hwy NE, Mailstop F17,
Atlanta, GA 30341, USA
E-mail: zca2@cdc.gov
Tel.: +1-770-4887982
Fax: +1-770-4884609

D. Hurtz III
Battelle Memorial Institute, Edgewood Operations,
2012 Tollgate Road, Suite 206, Bel Air, MD 21015, USA