Colorings of simplicial complexes and vector bundles over Davis–Januszkiewicz spaces

Dietrich Notbohm

Received: 20 May 2009 / Accepted: 23 June 2009 / Published online: 5 August 2009
© The Author(s) 2009. This article is published with open access at Springerlink.com

Abstract We show that coloring properties of a simplicial complex K are reflected by splitting properties of a bundle over the associated Davis–Januszkiewicz space whose Chern classes are given by the elementary symmetric polynomials in the generators of the Stanley–Reisner algebra of K .

Keywords Davis–Januszkiewicz spaces · Vector bundle · Characteristic classes · Colorings · Simplicial complexes

Mathematics Subject Classification (2000) 55R10 · 57R22 · 05C15

1 Introduction

For a simplicial complex K , Davis and Januszkiewicz constructed a family of spaces, all of which are homotopy equivalent, and whose integral cohomology is isomorphic to the associated Stanley–Reisner algebra $\mathbb{Z}[K]$ [2, Sect. 4]. We denote a generic model for this homotopy type by $DJ(K)$. In the above mentioned influential paper, Davis and Januszkiewicz also constructed a particular complex vector bundle λ over $DJ(K)$ whose Chern classes are given by the elementary symmetric polynomials in the generators of $\mathbb{Z}[K]$ [2, Sect. 6]. This vector bundle is of particular interest. For example, if K is the dual of the boundary of a simple polytope P , then the associated moment angle complex Z_K is a manifold and the realification $\lambda_{\mathbb{R}}$ of λ is stably isomorphic to the bundle given by applying the Borel construction to the tangent bundle of Z_K . And if M^{2n} is a quasitoric manifold over P , then again the Borel construction applied to the tangent bundle of M^{2n} produces a vector bundle stably isomorphic to $\lambda_{\mathbb{R}}$ [2, Theorem 6.6, Lemma 6.5].

D. Notbohm (✉)
Department of Mathematics, Faculty of Sciences, Vrije Universiteit Amsterdam,
De Boelelaan 1081a, 1081 HV Amsterdam, The Netherlands
e-mail: notbohm@few.vu.nl

Davis and Januszkiewicz also noticed that, if K is the dual of the boundary of a polytope of dimension n and admits a coloring with n colors, the bundle λ splits into a direct sum of n complex line bundles and a trivial bundle [2, Sect. 6.2]. We are interested in generalizations of this observation. In fact, we will show that a simplicial complex admits a coloring with r colors precisely when λ splits stably into a direct sum of r linear complex bundles and a trivial bundle. We will also show that a similar result holds for the realification of λ .

To make our statements more precise we have to fix notation and recall some basic constructions. Let $[m] := \{1, \dots, m\}$ be the set of the first m natural numbers. A finite abstract simplicial complex K on $[m]$ is given by a set of faces $\alpha \subseteq [m]$ which is closed under the formation of subsets. We consider the empty set $\emptyset$ as a face of K . The dimension $\dim \alpha$ of a face α is given in terms of its cardinality by $|\alpha| - 1$, and the dimension $\dim K$ of K is the maximum of the dimensions of its faces. The most basic examples are given by full simplices. For $\alpha \subseteq [m]$ we denote by $\Delta[\alpha]$ the simplicial complex which consists of all possible subsets of α . Then $\Delta[\alpha]$ is an $(|\alpha| - 1)$ -dimensional simplex. The full simplex $\Delta[m]$ contains K as a subcomplex, and if $\alpha \in K$ then $\Delta[\alpha] \subset K$ is a subcomplex as well.

A *regular r -paint coloring*, an *r -coloring* for short, of a simplicial complex K is a non degenerate simplicial map $g: K \rightarrow \Delta[r]$, i.e. g maps each face of K isomorphically on a face of $\Delta[r]$. The inclusion $K \subset \Delta[m]$ always provides an m -coloring. If $\dim(K) = n - 1$, then K may only allow r -colorings for $r \geq n$.

For a commutative ring R with unit we denote by $R[m] := R[v_1, \dots, v_m]$ the graded polynomial algebra generated by the algebraically independent elements $v_1, \dots, v_m$ of degree two, one for each vertex of K . For each subset $\alpha \subseteq [m]$ we denote by $v_\alpha := \prod_{j \in \alpha} v_j$ the square-free monomial whose factors are in one-to-one correspondence with vertices contained in α . The graded Stanley-Reisner algebra $R[K]$ associated with K is defined as the quotient $R[K] := R[m]/I_K$, where $I_K \subset R[m]$ is the ideal generated by all elements v_μ such that $\mu \subseteq [m]$ is not a face of K .

Since BT^m is an Eilenberg-MacLane space realizing the polynomial algebra $\mathbb{Z}[m]$, the projection $\mathbb{Z}[m] \rightarrow \mathbb{Z}[K]$ can be realized by a map $f: DJ(K) \rightarrow BT^m$. We can think of T^m as the maximal torus of the unitary group $U(m)$. The pull back along the composition $DJ(K) \rightarrow BT^m \rightarrow BU(m)$ of the universal bundle over $BU(m)$ gives a vector bundle $\lambda \downarrow DJ(K)$. This is the vector bundle studied by Davis and Januszkiewicz and mentioned above. The total Chern class $c(\lambda) = 1 + \sum_i c_i(\lambda)$ of λ is then given by $c(\lambda) = \prod_{i=1}^m (1 + v_i) \in \mathbb{Z}[K]$.

The realification of a complex vector bundle ξ is denoted by $\xi_{\mathbb{R}}$. Confusing notation we will denote by $\mathbb{C}$ and $\mathbb{R}$ a 1-dimensional trivial complex or real vector bundle over a space X . Now we can state our main theorem.

Theorem 1.1 *Let K be an finite simplicial complex over the vertex set $[m]$. Then the following conditions are equivalent.*

- (i) K admits an r -coloring $K \rightarrow \Delta[r]$.
- (ii) The vector bundle λ splits into a direct sum $(\bigoplus_{i=1}^r v_i) \oplus \mathbb{C}^{m-r}$ of r complex line bundles v_i and a trivial $(m - r)$ -dimensional complex bundle.
- (iii) The realification $\lambda_{\mathbb{R}}$ of λ splits into a sum $(\bigoplus_{i=1}^r \theta_i) \oplus \mathbb{R}^{2(m-r)}$ of r 2-dimensional real bundles θ_i and a trivial $2(m - r)$ -dimensional real bundle.
- (iv) The vector bundle λ is stably isomorphic to a direct sum $\bigoplus_{i=1}^r v_i$ of r complex line bundles.
- (v) The realification $\lambda_{\mathbb{R}}$ is stably isomorphic to a direct sum $\bigoplus_{i=1}^r \theta_i$ of r 2-dimensional real bundles.

Several of our vector bundles will be constructed as homotopy orbit spaces. For a compact Lie group G and a G -space X , the *Borel construction* or *homotopy orbit space* $EG \times_G X$ will be denoted by X_{hG} . If $\eta \downarrow X$ is an n -dimensional G -vector bundle over X with total space $E(\eta)$, the Borel construction establishes a fibre bundle $E(\eta)_{hG} \rightarrow X_{hG}$. In fact, this is an n -dimensional vector bundle over X_{hG} , denoted by η_{hG} . For definitions and details see [4].

Let M^{2n} be a quasitoric manifold over the simple polytope P . That is that M^{2n} carries a T^n -action, which is locally standard and that the orbit space $M^{2n}/T^n = P$ is a simple polytope. The Borel construction produces a space $(M^{2n})_{hT^n}$, which is homotopy equivalent to $DJ(K_P)$, where K_P is the simplicial complex dual to the boundary of P . For details see [2, Sect. 4.2]. Let τ_M denote the tangent bundle of M^{2n} . Davis and Januszkiewicz showed that the vector bundle $(\tau_M)_{hT^n} \downarrow DJ(K)$ and $\lambda_{\mathbb{R}}$ are stably isomorphic as real vector bundles over $DJ(K_P)$ [2, Sect. 6]. We can draw the following corollary of Theorem 1.1.

Corollary 1.2 *Let M^{2n} be a quasitoric manifold over a simple polytope P . Let K_P be the simplicial complex dual to the boundary of P . If the tangent bundle τ_M of M^{2n} is stably equivariantly isomorphic to a direct sum of r 2-dimensional real equivariant T^n -bundles over M^{2n} , then K_P admits an r -coloring.*

The paper is organized as follows. For the proof of our main theorem we will need two different models for $DJ(K)$. They are discussed in the next section. In Sect. 3, we will use some geometric constructions to produce a splitting of λ from a given coloring. The final section contains the proof of Theorem 1.1.

If not specified otherwise, K will always denote an $(n - 1)$ -dimensional finite simplicial complex with m -vertices.

2 Models for $DJ(K)$

Let $\text{CAT}(K)$ denote the category whose objects are the faces of K and whose arrows are given by the subset relations between the faces. $\text{CAT}(K)$ has an initial object given by the empty face. Given a pair (X, Y) of pointed topological space we can define covariant functors

$$X^K, (X, Y)^K : \text{CAT}(K) \rightarrow \text{TOP}.$$

The functor X^K assigns to each face α the cartesian product X^α and to each morphism $i_{\alpha, \beta}$ the inclusion $X^\alpha \subset X^\beta$ where missing coordinates are set to the base point $*$. If $\alpha = \emptyset$, then X^α is a point. And $(X, Y)^K$ assigns to α the product $X^\alpha \times Y^{[m] \setminus \alpha}$ and to $i_{\alpha, \beta}$ the coordinate wise inclusion $X^\alpha \times Y^{[m] \setminus \alpha} \subset X^\beta \times Y^{[m] \setminus \beta}$. The inclusions $X^\alpha \subset X^{[m]} = X^m$ and $X^\alpha \times Y^{[m] \setminus \alpha} \subset X^m$ establish inclusions

$$\text{colim}_{\text{CAT}(K)} X^K \rightarrow X^m, \quad \text{colim}_{\text{CAT}(K)} (X, Y)^K \rightarrow X^m.$$

We are interested in two particular cases, namely the functor X^K for the classifying space $BT = \mathbb{C}P^\infty$ of the 1-dimensional circle T and the functor $(X, Y)^K$ for the pair (D^2, S^1) . The colimit

$$Z_K := \text{colim}_{\text{CAT}(K)} (D^2, S^1)^K$$

is called the *moment angle complex* associated to K . The inclusions $Z_K \subset (D^2)^m \subset \mathbb{C}^m$ allow to restrict the standard T^m -action on $\mathbb{C}^m$ to Z_K . The Borel construction produces a fibration

$$q_K : (Z_K)_{hT^m} \rightarrow BT^m$$

with fiber Z_K . Moreover, $B_T K := (Z_K)_{hT^m}$ is a realization of the Stanley-Reisner algebra $\mathbb{Z}[K]$ and a model for $DJ(K)$. That is there exists an isomorphism $H^*(B_T K; \mathbb{Z}) \cong \mathbb{Z}[K]$ such that the map $H^*(q_K; \mathbb{Z})$ can be identified with the map $\mathbb{Z}[m] \rightarrow \mathbb{Z}[K]$ [2, Theorem 4.8]. We will use this model for geometric construction with our vector bundles.

Buchstaber and Panov gave a different construction for $DJ(K)$. They showed that $c(K) := \text{colim}_{\text{CAT}(K)} BT^K$ is homotopy equivalent to $B_T K$ and that the inclusion

$$c(K) \rightarrow BT^m$$

is homotopic to q_K [1, Theorem 6.29]. In particular, each face $\alpha \in K$ defines a map $h_\alpha: BT^\alpha \rightarrow c(K)$. The model $c(K)$ will be used to produce a coloring from a stable splitting of λ .

Remark 2.1 If K is the triangulation of an $(n - 1)$ -dimensional sphere, the moment angle complex Z_K is a manifold. In this case, the tangent bundle τ_Z is a $(m + n)$ -dimensional T^m -equivariant vector bundle, which satisfies the analogue of Corollary 1.2. If τ_Z is stably equivariantly isomorphic to a direct sum of r 2-dimensional real equivariant T^m -bundles over Z_K , then K admits an r coloring. Again this follows from the fact that $(\tau_Z)_{hT^m}$ and $\lambda_{\mathbb{R}}$ are stably isomorphic [2, Sect. 6].

3 Geometric constructions

The m -dimensional torus T^m acts coordinate wise on $\mathbb{C}^m$. And the diagonal action of T^m on $\mathbb{C}^m \times Z_K$ makes the projection $\mathbb{C}^m \times Z_K \rightarrow Z_K$ onto the second factor into a T^m -equivariant complex vector bundle over Z_K , denoted by λ' . An application of the Borel construction produces the bundle $\lambda := \lambda'_{hT^m} \downarrow B_T K$ over $B_T K$ whose total Chern class is given by $c(\lambda) = \prod_i (1 + v_i) \in \mathbb{Z}[K]$ and whose classifying map is the composition $B_T K \xrightarrow{q_K} BT^m \rightarrow BU(m)$. Since T^m acts coordinatewise on $\mathbb{C}^m$, both bundles, λ' and λ split into a direct sum of (equivariant) line bundles. Let $\mathbb{C}_j$ denote the j th component of $\mathbb{C}^m$. In particular, T^m acts on $\mathbb{C}_j$ via the projection $T^m \rightarrow T^{\{j\}}$ onto the j th component of T^m . The vector bundle $\lambda'_j := \mathbb{C}_j \times Z_K$ is T^m -equivariant, and $\lambda_j := (\lambda'_j)_{hT^m}$ is a 1-dimensional complex vector bundle over $B_T K$. We have $\lambda' \cong \bigoplus_j \lambda'_j$ and $\lambda \cong \bigoplus_j \lambda_j$. All this can be found in [2, Sect. 6].

If $g: K \rightarrow \Delta[r]$ is an r -coloring we want to construct an equivariant splitting of $\lambda' \downarrow Z_K$ into a direct sum of T^m -equivariant complex line bundles and a trivial bundle $\mathbb{C}^{m-r}$. We will use ideas of Davis and Januszkiewicz discussed in [2, Sect. 6.2]. For each $i \in [r]$ we denote by $S_i := g^{-1}(i) \subset [m]$ the preimage of i and by $s_i := |S_i|$ the order of S_i . There are two vector bundles associated with S_i , namely the tensor product $v_i := \bigotimes_{j \in S_i} \lambda'_j$ of all complex line bundles associated to the vertices contained in S_i and the direct sum $\eta_i := \bigoplus_{j \in S_i} \lambda'_j$ of all these line bundles. Both are T^m -equivariant vector bundles over Z_K .

Lemma 3.1 *For all $i \in [r]$, there exists an T^m -equivariant vector bundle isomorphism $v_i \oplus \mathbb{C}^{s_i-1} \rightarrow \eta_i$.*

For simplicial complexes dual to the boundary of simple polytopes the claim is already stated in [2, Sect. 6.2]. We will give here a different proof.

Proof For simplification we drop the subindex i in the notation and assume that $S = [s]$. We will think of $\mathbb{C}^{s-1} \subset \mathbb{C}^s$ as the subspace given by $\{(x_1, \dots, x_s) \in \mathbb{C}^s \mid \sum_k x_k = 0\}$. We define a map

$$f : \mathbb{C} \times \mathbb{C}^{s-1} \times Z_K \longrightarrow \mathbb{C}^s \times Z_K$$

by $f(y, x, z) := (u, z)$ where the j th coordinate u_j of u is given by $u_j := y \prod_{k \neq j, k \in [s]} \bar{z}_k + z_j x_j$. Here, $\bar{z}_k$ denotes the complex conjugate of z_k . If T^m acts on $\mathbb{C}$ via the map $t \mapsto \prod_{j \in [s]} t_j$, trivially on $\mathbb{C}^{s-1}$ and on $\mathbb{C}^s$ via the projection $T^m \longrightarrow T^s$ onto the first s coordinates, one can easily show that this map is T^m -equivariant. Moreover, with these actions, the source is the total space of the bundle $\nu \oplus \mathbb{C}^{s-1} \downarrow Z_K$ and the target the total space of $\eta \downarrow Z_K$. Since both sides have the same dimension, it is only left to show that f is fiber wise a monomorphism.

By construction, any subset $\{j, k\} \subset [s]$ is a missing face in K . Since $Z_K = \bigcup_{\alpha \in K} (D^2)^\alpha \times (S^1)^{[m] \setminus \alpha}$, the space $(D^2)^{\{j, k\}} \times (S^1)^{[m] \setminus \{i, j\}}$ is not contained in Z_K and for $z = (z_1, \dots, z_m) \in Z_K$ there is at most one coordinate among $z_1, \dots, z_s$ which is trivial.

Now we assume that $f(y, x, z) = (0, z)$. In particular, we have $x_j z_j = -y \prod_{k \neq j} \bar{z}_k$. If one of the coordinates z_j vanishes, say $z_1 = 0$, then $z_j \neq 0$ for $j \neq 1$ and hence $y = 0$ as well as $x_j = 0$ for $j \neq 1$. Since $\sum_j x_j = 0$, we also have $x_1 = 0$.

If $z_j \neq 0$ for all j , then $x_j = y \prod_{k \neq j} \bar{z}_k / z_j$ and $0 = \sum_j x_j = \sum_j y \prod_{k \neq j} \bar{z}_k / z_j = y \sum_j \prod_{k \neq j} \bar{z}_k / z_j$. Multiplying with $\prod_j z_j$ shows that $y \sum_j \prod_{k \neq j} \bar{z}_k z_k = 0$ and hence that $y = 0$ as well as $x_j = 0$ for all j . This shows that f is a fiber wise monomorphism and finishes the proof. $\square$

Corollary 3.2 *Let $K \longrightarrow \Delta[r]$ be an r -coloring of a finite simplicial complex. Then the following holds:*

- (i) *The bundle $\lambda' \downarrow Z_K$ splits equivariantly into a direct sum of r equivariant complex line bundles and a trivial bundle.*
- (ii) *The bundle $\lambda \downarrow DJ(K)$ splits into a direct sum of r complex line bundles and a trivial bundle.*

Proof By Proposition 3.1 we have

$$\lambda' \cong \bigoplus_{j=1}^r \bigoplus_{i \in S_j} \lambda'_i \cong \bigoplus_{j=1}^r (v'_j \oplus \mathbb{C}^{s_i-1}) \cong \left(\bigoplus_{j=1}^r v'_j \right) \oplus \mathbb{C}^{m-r}.$$

This proves the first part, the second follows from the first by applying the Borel construction. $\square$

4 Proof of Theorem 1.1

The proof needs some preparation. For topological spaces X and Y we denote by $[X, Y]$ the set of homotopy classes of maps from X to Y and for two compact Lie groups G and H by $\text{hom}(H, G)$ the set of Lie group homomorphism $H \longrightarrow G$.

Let G be a compact connected Lie group with maximal torus $j : T_G \hookrightarrow G$ and Weyl group W_G . Since the action of W_G on T_G is induced by conjugation with elements of G , the composition of $w \in W_G$ and j induces a map between the classifying spaces homotopic to Bj . And passing to classifying spaces followed by composing with Bj induces a map $\text{hom}(H, T_G) \longrightarrow [BH, BG]$ which factors through the orbit space of the W_G -action on $\text{hom}(H, T_G)$ and provides a map $\text{hom}(H, T_G)/W_G \longrightarrow [BH, BG]$. The following two facts may be found in [3] and are needed for the proof of our main theorem.

Theorem 4.1 [3] *Let G be a connected compact Lie group and S a torus.*

- (i) *The map $\text{hom}(S, T_G)/W_G \rightarrow [BS, BG]$ is a bijection.*
- (ii) *The map $[BS, BG] \rightarrow \text{Hom}(H^*(BG; \mathbb{Q}), H^*(BS; \mathbb{Q}))$ is an injection.*

The rational cohomology $H^*(BG; \mathbb{Q}) \cong H^*(BT_G, \mathbb{Q})^{W_G}$ is the ring of polynomial invariants of the induced W_G -action on the polynomial algebra $H^*(BT_G; \mathbb{Q})$. For $G = SO(2k+1)$ the maximal torus $T_{SO(2k+1)} = T^k$ is an k -dimensional torus and we can identify $H^*(BT_{SO(2k+1)}; \mathbb{Z})$ with $\mathbb{Z}[k] = \mathbb{Z}[v_1, \dots, v_k]$. The Weyl group $W_{SO(2k+1)}$ is the wreath product $\mathbb{Z}/2 \wr \Sigma_k$ where $(\mathbb{Z}/2)^k$ acts on T^k via coordinate wise complex conjugation and Σ_k via permutations of the coordinates. The rational cohomology of $BSO(2k+1)$ is then given by

$$H^*(BSO(2k+1); \mathbb{Q}) \cong \mathbb{Q}[k]^{\mathbb{Z}/2 \wr \Sigma_k} \cong \mathbb{Q}[p_1, \dots, p_k].$$

The classes p_i are already defined over $\mathbb{Z}$. On the one hand $p_i \in H^{4i}(BSO(2k+1); \mathbb{Z})$ is the universal i th Pontrjagin class for oriented bundles and on the other hand $p_i = (-1)^i \sigma_i(v_1^2, \dots, v_k^2) \in \mathbb{Z}[k]^{\mathbb{Z}/2 \wr \Sigma_k}$ is up to a sign the i th elementary symmetric polynomial in the squares of the generators of $\mathbb{Z}[k]$. In particular, for an oriented $(2k+1)$ -dimensional real vector bundle ρ over a space X , the total Pontrjagin class $p(\rho) = 1 + \sum_{i=1}^k p_i(\rho)$ determines completely the map $H^*(BSO(2k+1); \mathbb{Q}) \rightarrow H^*(X; \mathbb{Q})$ induced by the classifying map $\rho: X \rightarrow BSO(2k+1)$.

Example 4.2 Let $\rho: BT^s \rightarrow BSO(2k+1)$ be the composition of a coordinate wise inclusion $\hat{\rho}: BT^s \rightarrow BT^k$ followed by the maximal torus inclusion $BT^k \rightarrow BSO(2k+1)$. Then the total Pontrjagin class of ρ is given by $p(\rho) = \prod_{i=1}^s (1 - v_i^2)$, where we identify $H^*(BT^s; \mathbb{Z})$ with $\mathbb{Z}[v_1, \dots, v_s]$.

By Theorem 4.1, up to homotopy every vector bundle $\omega: BT^s \rightarrow BSO(2k+1)$ is the composition of a lift $\hat{\omega}: BT^s \rightarrow BT^k$ and Bj . If $p(\omega) = p(\rho)$ then both maps ρ and ω are homotopic and the underlying homomorphisms $j_\omega, j_\rho: T^s \rightarrow T^k$ of the lifts $\hat{\omega}$ and $\hat{\rho}$ differ only by an element of the Weyl group (Theorem 4.1). In particular, since j_ρ is given by a coordinatewise inclusion, the homomorphism j_ω also is a coordinate wise inclusion possibly followed by complex conjugation on some coordinates.

Proof of Theorem 1.1 If $K \rightarrow \Delta[r]$ is an r -coloring, Corollary 3.2 provides the appropriate splitting for λ . The splitting conditions on λ can be put into a hierarchy, a splitting of λ establishes a splitting of $\lambda_{\mathbb{R}}$ and a stable isomorphism between λ and a direct sum of r complex line bundles, and the two latter conditions a stable isomorphism between $\lambda_{\mathbb{R}}$ and a direct sum of r 2-dimensional real bundles. It is only left to show that this last stable isomorphism allows to construct a coloring.

We will work with the model $c(K)$ for $DJ(K)$ and again describe vector bundles over $c(K)$ by their classifying maps. In particular, for $t \geq 2m$ the real vector bundle $\rho_t := \lambda_{\mathbb{R}} \oplus \mathbb{R}^{t-2m}$ is a map $\rho_t: c(K) \rightarrow BO(t)$. Since we are considering stable splittings, we can pass from ρ_t to ρ_{t+1} , if necessary, and assume that $t = 2s + 1$ is odd. This will simplify the discussion. For example, if t is odd, we have $BO(t) \simeq BSO(t) \times B\mathbb{Z}/2$. And since $c(K)$ is simply connected, the bundle ρ_t has a unique orientation given by the first coordinate of the map $\rho_t: c(K) \rightarrow BSO(t) \times B\mathbb{Z}/2$. We also denote this map by ρ_t . The total Pontrjagin class of ρ_t is given by $p(\rho_t) = \prod_{i=1}^m (1 - v_i^2)$ [2, Sect. 6].

Let $\phi: BT^r \rightarrow BSO(t)$ be the map induced by the composition of the coordinate wise inclusion $T^r \subset T^s$ into the first r coordinates followed by the maximal torus inclusion $T^s = T_{SO(t)} \subset SO(t)$. A splitting $\rho_t \cong (\bigoplus_{j=1}^r \theta_j) \oplus \mathbb{R}^{t-2r}$ establishes a map $\hat{\rho}_t: c(K) \rightarrow BT^r$ such that $\phi \hat{\rho}_t \simeq \rho_t$.

Now let $\alpha \in K$ be a face and $h_\alpha: BT^\alpha \rightarrow c(K)$ the associated map. The composition $\hat{\rho}_t h_\alpha: BT^\alpha \rightarrow BT^r$ determines a unique homomorphism $j_\alpha: T^\alpha \rightarrow T^r$. The total Pontrjagin class of $\phi \hat{\rho}_t h_\alpha$ is given by $p(\phi \hat{\rho}_t h_\alpha) = \prod_{i \in \alpha} (1 - v_i^2)$. Example 4.2 shows that j_α is a coordinate wise inclusion $T^\alpha \rightarrow T^k$ possibly followed by complex conjugation on some coordinates. The coordinate wise inclusion defines an injection $\alpha \rightarrow [r]$. Since for any inclusion of faces $\beta \subset \alpha$ the restriction $(\hat{\rho}_t h_\alpha)|_{BT^\beta}$ equals the composition $\hat{\rho}_t h_\beta$, the underlying homomorphisms satisfies the formula $j_\alpha|_\beta = j_\beta$. We can conclude that the collection of all these maps defines a map $[m] \rightarrow [r]$, whose restriction to any face of K is an injection. This establishes a non degenerate simplicial map $K \rightarrow \Delta[r]$ which is an r -coloring for K . $\square$

Acknowledgements We would like to thank Nigel Ray and Natalia Dobrinskaya for many helpful discussions.

Open Access This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

References

1. Buchstaber, V.M., Panov, T.E.: Torus actions and their applications in topology and combinatorics, vol. 24, University Lecture Series. American Mathematical Society (2002)
2. Davis, M.W., Januszkiewicz, T.: Convex Polytopes, Coxeter Orbifolds and Torus Actions. *Duke Math. J.* **62**, 417–451 (1991)
3. Notbohm, D.: Maps between classifying spaces. *Math. Z.* **207**, 153–168 (1991)
4. Segal, G.: Equivariant K -theory. *Publ. Math. Inst. Hautes Étud. Sci.* **34**, 129–151 (1968)