On splitting theorems for CAT(0) spaces and compact geodesic spaces of non-positive curvature

Tetsuya Hosaka

Received: 13 June 2011 / Accepted: 24 October 2011 / Published online: 13 December 2011
© The Author(s) 2011. This article is published with open access at Springerlink.com

Abstract In this paper, we show some splitting theorems for CAT(0) spaces on which a product group acts geometrically and we obtain a splitting theorem for compact geodesic spaces of non-positive curvature. A CAT(0) group Γ is said to be *rigid*, if Γ determines its boundary up to homeomorphisms of a CAT(0) space on which Γ acts geometrically. C. Croke and B. Kleiner have constructed a non-rigid CAT(0) group. As an application of the splitting theorems for CAT(0) spaces, we obtain that if Γ_1 and Γ_2 are rigid CAT(0) groups then so is $\Gamma_1 \times \Gamma_2$.

Keywords Splitting theorem · CAT(0) space · Non-positive curvature · Boundary · CAT(0) group · Rigid

Mathematics Subject Classification (2000) 20F65 · 57M07

1 Introduction and preliminaries

The purpose of this paper is to show some splitting theorems for CAT(0) spaces on which a product group acts geometrically (i.e. properly and cocompactly by isometries). As applications, we obtain a splitting theorem for compact geodesic spaces of non-positive curvature and rigidity of the product of rigid CAT(0) groups.

On splitting theorems for spaces of non-positive curvature, we can find historical research in [3, 7, 9, 12] and [10].

Partly supported by the Grant-in-Aid for Young Scientists (B), The Ministry of Education, Culture, Sports, Science and Technology, Japan. (No. 21740037).

T. Hosaka (✉)
Department of Mathematics, Shizuoka University,
Suruga-ku, Shizuoka 422-8529, Japan
e-mail: sthosak@ipc.shizuoka.ac.jp

After some preliminaries on CAT(0) spaces and their boundaries in Sect. 2, we first show the following splitting theorem by a similar argument to the proof of [3, Proposition II.6.23] in Sect. 3.

Theorem 1 *Suppose that a group $\Gamma = \Gamma_1 \times \Gamma_2$ acts geometrically on a CAT(0) space X . If Γ_1 acts cocompactly on the convex hull $C(\Gamma_1 x_0)$ of some Γ_1 -orbit, then there exists a closed, convex, Γ -invariant, quasi-dense subspace $X' \subset X$ such that X' splits as a product $X_1 \times X_2$, Γ_1 naturally acts geometrically on X_1 , and Γ_2 acts geometrically on X_2 by projection. Here each subspace of the form $X_1 \times \{x_2\}$ is the closed convex hull of some Γ_1 -orbit.*

Here we define the induced action by projection in Sect. 2.

From Theorem 1, we obtain the following splitting theorem which corresponds to [3, Theorem II.6.21]. (N. Monod [10, Corollary 10] independently has proved the generalization of this splitting theorem.)

Theorem 2 *Suppose that a group $\Gamma = \Gamma_1 \times \Gamma_2$ acts geometrically on a CAT(0) space X . If the center of Γ is finite, then there exists a closed, convex, Γ -invariant, quasi-dense subspace $X' \subset X$ such that X' splits as a product $X_1 \times X_2$ and the action of $\Gamma = \Gamma_1 \times \Gamma_2$ on $X' = X_1 \times X_2$ is the product action.*

Theorem 2 has a overlap with a result in [10].

In [12] and [10], some splitting theorems for Hadamard manifolds and CAT(0) spaces have been proved. In their splitting theorems [12, Theorem 1] and [10, Corollary 10] and also in Theorem 2, the assumption “there is not a Γ -fixed point in ∂X ” in [12, Theorem 1] and [10, Corollary 10] and the assumption “the center of Γ is finite” in Theorem 2 are very important, and in these cases, the action does split. Here we note that in the case a group Γ acts geometrically on a CAT(0) space X , Γ has finite center if and only if there is not a Γ -fixed point in ∂X (cf. Lemma 2.4).

For example, it is known that every Coxeter group is a CAT(0) group whose center is finite.

In Sect. 4, we prove the following splitting theorem which is the main theorem in this paper and in which we do not need the assumption that “there is not a Γ -fixed point in ∂X ” and that “the center of Γ is finite”.

Theorem 3 *Suppose that a group $\Gamma = \Gamma_1 \times \Gamma_2$ acts geometrically on a CAT(0) space X . Then there exist finite-index subgroups G_1 and G_2 of Γ_1 and Γ_2 respectively and there exist closed convex subspaces X_1, X_2, X'_1, X'_2 in X such that*

- (1) $X_1 \times X'_2$ is a closed convex $\Gamma_1 \times G_2$ -invariant quasi-dense subspace of X ,
- (2) $X'_1 \times X_2$ is a closed convex $G_1 \times \Gamma_2$ -invariant quasi-dense subspace of X ,
- (3) X'_1 and X'_2 are isometric to some quasi-dense subspaces of X_1 and X_2 respectively,
- (4) there exist geometric actions of Γ_1 and Γ_2 on X_1 and X_2 respectively, and
- (5) there exist geometric actions of G_1 and G_2 on X'_1 and X'_2 respectively.

Hence the boundary ∂X of X is homeomorphic to the join $\partial X_1 * \partial X_2$ of the boundaries of X_1 and X_2 .

Theorem 3 does not have a overlap with the results in [10].

In this splitting theorem, the center of Γ need not be finite and also we do not need the assumption that there is not a Γ -fixed point in ∂X . On the other hand, the action does not split in general. For example, the geometric action of $\Gamma = \mathbb{Z} \times \mathbb{Z}$ on $X = \mathbb{R} \times \mathbb{R}$ defined by $(a, b) \cdot (x, y) = (a + x, a + b + y)$ for $(a, b) \in \mathbb{Z} \times \mathbb{Z}$ and $(x, y) \in \mathbb{R} \times \mathbb{R}$ does not split (of course, the CAT(0) space X splits).

A CAT(0) space X is said to have the *geodesic extension property*, if every geodesic can be extended to a geodesic line $\mathbb{R} \rightarrow X$. On CAT(0) spaces with the geodesic extension property, we obtain the following splitting theorem as an application of Theorem 3.

Theorem 4 *Suppose that a group $\Gamma = \Gamma_1 \times \Gamma_2$ acts geometrically on a CAT(0) space X with the geodesic extension property. Then X splits as a product $X_1 \times X_2$ and there exist geometric actions of Γ_1 and Γ_2 on X_1 and X_2 , respectively. Moreover if Γ has finite center, then Γ preserves the splitting, i.e., the action of $\Gamma = \Gamma_1 \times \Gamma_2$ on $X = X_1 \times X_2$ is the product action.*

In Sect. 5, we study products of CAT(0) groups. A group Γ is called a *CAT(0) group*, if Γ acts geometrically on some CAT(0) space.

Theorem 3 implies the following.

Theorem 5 *Γ_1 and Γ_2 are CAT(0) groups if and only if $\Gamma_1 \times \Gamma_2$ is a CAT(0) group.*

In [5], C. Croke and B. Kleiner have proved that there exists a CAT(0) group Γ and CAT(0) spaces X and Y such that Γ acts geometrically on X and Y and the boundaries of X and Y are not homeomorphic. On CAT(0) groups and their boundaries, it is an important open problem when does a CAT(0) group determine its boundary up to homeomorphisms. A CAT(0) group Γ is said to be *rigid*, if Γ determines its boundary up to homeomorphisms of a CAT(0) space on which Γ acts geometrically. Then we denote the boundary of the rigid CAT(0) group Γ by $\partial\Gamma$.

A conclusion in [3, Theorem II.7.1] and [2] implies that if Γ is a rigid CAT(0) group, then $\Gamma \times \mathbb{Z}^n$ is also a rigid CAT(0) group for each $n \in \mathbb{N}$. In [11], K. Ruane has proved that if $\Gamma_1 \times \Gamma_2$ is a CAT(0) group and if Γ_1 and Γ_2 are hyperbolic groups (in the sense of Gromov) then $\Gamma_1 \times \Gamma_2$ is rigid. On rigidity of products of rigid CAT(0) groups, we obtain the following theorem from Theorem 3 as a generalization of these results.

Theorem 6 *If Γ_1 and Γ_2 are rigid CAT(0) groups, then so is $\Gamma_1 \times \Gamma_2$, and the boundary $\partial(\Gamma_1 \times \Gamma_2)$ is homeomorphic to the join $\partial\Gamma_1 * \partial\Gamma_2$ of the boundaries of Γ_1 and Γ_2 .*

2 Lemmas on CAT(0) spaces and their boundaries

In this section, we introduce some properties of CAT(0) spaces and their boundaries and show some lemmas needed later.

Definitions and details of CAT(0) spaces and their boundaries are found in [3] and [6]. A *geometric action* on a CAT(0) space is an action by isometries which is proper ([3, p.131]) and cocompact. We note that every CAT(0) space on which some group acts geometrically is a proper space ([3, p.132]).

Let X be a metric space and let γ be an isometry of X . Then the *translation length* of γ is defined as $|\gamma| = \inf\{d(x, \gamma x) \mid x \in X\}$, and the *minimal set* of γ is defined as $\text{Min}(\gamma) = \{x \in X \mid d(x, \gamma x) = |\gamma|\}$. If Γ is a group acting by isometries on X , then $\text{Min}(\Gamma) := \bigcap_{\gamma \in \Gamma} \text{Min}(\gamma)$. An isometry γ is said to be *semi-simple* if $\text{Min}(\gamma)$ is non-empty.

Also an isometry γ of a metric space X is called

- (1) *elliptic* if γ has a fixed point,
- (2) *hyperbolic* if γ is semi-simple and not elliptic, and
- (3) *parabolic* if γ is not semi-simple.

Now we define induced actions by projection.

Definition 2.1 Suppose that a group $\Gamma = \Gamma_1 \times \Gamma_2$ acts geometrically on a CAT(0) space $X = X_1 \times X_2$. In general, the action does not split. Let $p_1 : X_1 \times X_2 \rightarrow X_1$ be the projection. We can define an action “ $*$ ” of Γ_1 on X_1 by $\gamma_1 * x_1 = p_1(\gamma_1 \cdot x_1)$ for $\gamma_1 \in \Gamma_1$ and $x_1 \in X_1$ (cf. [2]), where “ $\cdot$ ” is the original action of Γ on X . Then we say that the action “ $*$ ” of Γ_1 on X_1 is the induced action by the projection of the action “ $\cdot$ ”.

In this paper, we often say that Γ_1 *acts geometrically on X_1 by projection*, if the induced action of Γ_1 on X_1 by the projection of the original action of Γ on X is geometric.

Also, in this paper, we often say that Γ_1 *acts geometrically on X_1 by restriction*, if Γ_1 acts geometrically on X_1 by the restriction of the original action of Γ on X (hence, in such a case, X_1 is Γ_1 -invariant).

The following theorem is called the Flat Torus Theorem (cf. [1], [2, Proposition 1.1], [3, Theorem II.7.1]).

Theorem 2.2 ([3, Theorem II.7.1]) *Let G be a group and let A be a free abelian group of rank n . Suppose that $\Gamma = G \times A$ acts geometrically on a CAT(0) space X . Then*

- (1) $\text{Min}(A) = \bigcap_{\alpha \in A} \text{Min}(\alpha)$ is a closed, convex, Γ -invariant, quasi-dense subspace of X that splits as a product $Y \times Z$, where Z is isometric to $\mathbb{R}^n$,
- (2) G acts geometrically on Y by projection, and
- (3) A acts geometrically on Z by restriction (moreover, Z is the convex hull of some orbit Ax_0 of X).

Here a subset X' of a metric space X is said to be *quasi-dense* if there exists a number $N > 0$ such that each point of X is N -close to some point of X' , i.e., $B(X', N) = X$.

Some results on “ Γ -invariant quasi-dense subspace” in this paper relate to [1, Corollary 2.7] and [4].

We show some lemmas needed later.

Lemma 2.3 *Let γ be a parabolic isometry of a proper cocompact CAT(0) space X . Then there exists a non-fixed point $\alpha \in \partial X$ of γ .*

Proof Let γ be a parabolic isometry of a proper cocompact CAT(0) space X . Then the isometry γ of X induces a homeomorphism of the boundary ∂X (cf. [3, Corollary II.8.9]). We suppose that the induced homeomorphism $\gamma : \partial X \rightarrow \partial X$ is the identity.

Since γ is a parabolic isometry, $\text{Min}(\gamma)$ is the empty set. Hence X is non-compact and ∂X is non-empty. This means that there exists a geodesic ray in X . Since X is cocompact, there exists a geodesic line $\sigma_1 : \mathbb{R} \rightarrow X$ by [3, Lemma II.9.34]. Let Y_1 be the union of the images of all geodesic lines parallel to σ_1 . For each geodesic line τ parallel to σ_1 , by hypothesis, $\gamma\tau(\infty) = \tau(\infty)$ and $\gamma\tau(-\infty) = \tau(-\infty)$. Hence $\gamma\tau$ is also parallel to σ_1 by the Flat Strip Theorem [3, Theorem II.2.13]. Thus Y_1 is γ -invariant. By [3, Theorem II.2.14], Y_1 is isometric to $X_1 \times \mathbb{R}$ for some convex subspace $X_1 \subset Y_1$. Then the restriction of γ to $Y_1 = X_1 \times \mathbb{R}$ splits as (γ_1, γ'_1) , where γ_1 and γ'_1 are isometries of X_1 and $\mathbb{R}$, respectively. Since γ'_1 is semi-simple (cf. [3, Proposition II.6.5]) and γ is not semi-simple, by [3, Proposition II.6.9], γ_1 is not semi-simple, i.e., γ_1 is a parabolic isometry. We show that the induced homeomorphism $\gamma_1 : \partial X_1 \rightarrow \partial X_1$ is the identity. Let ξ be a geodesic ray in X_1 . Then $\gamma\xi$ is a geodesic ray in γX_1 and $\gamma\xi(\infty) = \xi(\infty)$ because γ is the identity of ∂X . Hence ξ and $\gamma\xi$ are parallel and $\gamma_1\xi(\infty) = \xi(\infty)$ by the definition of γ_1 (since γ_1 is the projection on X_1 of γ). Thus $\gamma_1 : \partial X_1 \rightarrow \partial X_1$ is the identity.

Since γ_1 is a parabolic isometry, X_1 is non-compact and ∂X_1 is non-empty, hence there exists a geodesic ray ξ_1 in X_1 . Then

$$X \supset Y_1 = X_1 \times \mathbb{R} \supset \text{Im } \xi_1 \times \mathbb{R}.$$

Since X is cocompact, there exists a subspace Z of X which is isometric to $\mathbb{R}^2$ by [3, Lemma II.9.34]. Here we redefine σ_1 as a geodesic line in Z , and we also redefine Y_1 , X_1 and γ_1 from σ_1 . Then there exists a geodesic line σ_2 in X_1 . Indeed we can construct σ_2 as $\text{Im } \sigma_2 = Z \cap X_1$. Let Y_2 be the union of the images of all geodesic lines parallel to σ_2 in X_1 . Here Y_2 is γ_1 -invariant and isometric to $X_2 \times \mathbb{R}$ for some convex subspace X_2 of Y_2 by [3, Theorem II.2.14]. Then the restriction of γ_1 to $Y_2 = X_2 \times \mathbb{R}$ splits as (γ_2, γ'_2) , γ_2 is a parabolic isometry of X_2 and the induced homeomorphism γ_2 of ∂X_2 is the identity.

By iterating this argument, for any $n \in \mathbb{N}$, there exists a sequence

$$\begin{aligned} X \supset Y_1 &= X_1 \times \mathbb{R} \supset Y_2 \times \mathbb{R} = X_2 \times \mathbb{R}^2 \\ &\supset Y_3 \times \mathbb{R}^2 = X_3 \times \mathbb{R}^3 \supset \cdots \supset X_n \times \mathbb{R}^n, \end{aligned}$$

that is, $\mathbb{R}^n \subset X$ for any $n \in \mathbb{N}$. However this contradicts [3, Lemma II.7.4]. Thus $\gamma : \partial X \rightarrow \partial X$ is not the identity, i.e., $\gamma\alpha \neq \alpha$ for some $\alpha \in \partial X$. $\square$

Lemma 2.4 *Suppose that a group Γ acts geometrically on a CAT(0) space X . The center of Γ is finite if and only if there does not exist $\alpha \in \partial X$ such that $\Gamma\alpha = \alpha$.*

Proof We first show that if the center of Γ is finite then there does not exist $\alpha \in \partial X$ such that $\Gamma\alpha = \alpha$.

Suppose that there exists $\alpha \in \partial X$ such that $\Gamma\alpha = \alpha$. Then we show that Γ has infinite center. Let $x_0 \in X$ and let $\xi : [0, \infty) \rightarrow X$ be the geodesic ray such that $\xi(0) = x_0$ and $\xi(\infty) = \alpha$. We note that $\delta\xi(\infty) = \xi(\infty)$ for any $\delta \in \Gamma$ because $\Gamma\alpha = \alpha$. Since the action of Γ on X is cocompact, $\Gamma B(x_0, N) = X$ for some $N > 0$. Then there exists a sequence $\{\gamma_i \mid i \in \mathbb{N}\} \subset \Gamma$ such that $d(\gamma_i x_0, \text{Im } \xi) \leq N$ and $\gamma_i \neq \gamma_j$ if $i \neq j$.

Now we show that for each $\delta \in \Gamma$, there exists a subsequence $\{\gamma_{i_j} \mid j \in \mathbb{N}\} \subset \{\gamma_i \mid i \in \mathbb{N}\}$ such that $(\gamma_{i_1} \gamma_{i_j}^{-1})\delta = \delta(\gamma_{i_1} \gamma_{i_j}^{-1})$ for any $j \in \mathbb{N}$.

Let $\delta \in \Gamma$. Since $\delta\xi(\infty) = \xi(\infty)$, there exists a number $K_\delta > 0$ such that $d(\xi(r), \delta\xi(r)) \leq K_\delta$ for any $r \geq 0$. For each $i \in \mathbb{N}$, $d(\gamma_i x_0, \xi(r_i)) \leq N$ for some $r_i \geq 0$ because $d(\gamma_i x_0, \text{Im } \xi) \leq N$. Then

$$\begin{aligned} d(x_0, \gamma_i^{-1} \delta \gamma_i x_0) &= d(\gamma_i x_0, \delta \gamma_i x_0) \\ &\leq d(\gamma_i x_0, \xi(r_i)) + d(\xi(r_i), \delta\xi(r_i)) + d(\delta\xi(r_i), \delta \gamma_i x_0) \\ &\leq N + K_\delta + d(\xi(r_i), \gamma_i x_0) \\ &\leq 2N + K_\delta. \end{aligned}$$

Since the action of Γ on X is proper, $\{\gamma \in \Gamma \mid d(x_0, \gamma x_0) \leq 2N + K_\delta\}$ is a finite set. Hence there exists a subsequence $\{\gamma_{i_j} \mid j \in \mathbb{N}\} \subset \{\gamma_i \mid i \in \mathbb{N}\}$ such that $\gamma_{i_j}^{-1} \delta \gamma_{i_j} = \gamma_{i_1}^{-1} \delta \gamma_{i_1}$ for any $j \in \mathbb{N}$. Then $(\gamma_{i_1} \gamma_{i_j}^{-1})\delta = \delta(\gamma_{i_1} \gamma_{i_j}^{-1})$ for any $j \in \mathbb{N}$.

The CAT(0) group Γ is finitely presented (cf. [3, Theorem III.1.1(1)]). Let $\{\delta_1, \dots, \delta_n\}$ be a generating set of Γ . Then by iterating the above argument, there exists a subsequence $\{g_i \mid i \in \mathbb{N}\} \subset \{\gamma_i \mid i \in \mathbb{N}\}$ such that $(g_1 g_i^{-1})\delta_k = \delta_k(g_1 g_i^{-1})$ for any $i \in \mathbb{N}$ and $k \in \{1, \dots, n\}$. Since $\{\delta_1, \dots, \delta_n\}$ generates Γ ,

$$(g_1 g_i^{-1})\delta = \delta(g_1 g_i^{-1})$$

for any $i \in \mathbb{N}$ and $\delta \in \Gamma$. Here we note that $g_1 g_i^{-1} \neq g_1 g_j^{-1}$ if $i \neq j$ by the construction of $\{\gamma_i \mid i \in \mathbb{N}\}$. Thus the center of Γ contains the infinite set $\{g_1 g_i^{-1} \mid i \in \mathbb{N}\}$.

Next we show that if there does not exist $\alpha \in \partial X$ such that $\Gamma\alpha = \alpha$, then the center of Γ is finite.

Suppose that the center $Z(\Gamma)$ of Γ is infinite. Since the set $\{\delta x_0 \mid \delta \in Z(\Gamma)\}$ is unbounded in X , there exists a sequence

$$\{\delta_i x_0 \mid i \in \mathbb{N}\} \subset \{\delta x_0 \mid \delta \in Z(\Gamma)\}$$

which converges to some point $\alpha_0 \in \partial X$ in $X \cup \partial X$. Let $\gamma \in \Gamma$. Then the sequence $\{\gamma \delta_i x_0 \mid i \in \mathbb{N}\}$ converges to $\gamma \alpha_0$. Here since $\gamma \delta_i = \delta_i \gamma$,

$$\{\gamma \delta_i x_0 \mid i \in \mathbb{N}\} = \{\delta_i \gamma x_0 \mid i \in \mathbb{N}\} = \{\delta_i y_0 \mid i \in \mathbb{N}\},$$

where $y_0 = \gamma x_0$, which converges to α_0 in $X \cup \partial X$. Hence $\gamma \alpha_0 = \alpha_0$ for any $\gamma \in \Gamma$. Thus $\Gamma \alpha_0 = \alpha_0$. $\square$

3 Some splitting theorems

We first generalize the splitting theorem [3, Proposition II.6.23] as follows. This plays a key role to prove the main results.

Theorem 3.1 *Suppose that a group $\Gamma = \Gamma_1 \times \Gamma_2$ acts geometrically on a CAT(0) space X . If Γ_1 acts cocompactly on the convex hull $C(\Gamma_1 x_0)$ of some Γ_1 -orbit, then there exists a closed, convex, Γ -invariant, quasi-dense subspace $X' \subset X$ such that*

- (1) X' splits as a product $X_1 \times X_2$,
- (2) Γ_1 acts geometrically on X_1 by restriction (moreover, each subspace of the form $X_1 \times \{x_2\}$ is the closed convex hull of some Γ_1 -orbit), and
- (3) Γ_2 acts geometrically on X_2 by projection.

Proof We show this theorem by a similar proof to the one in [3, pp. 240–241].

Let Σ be the set of closed, convex, non-empty, Γ_1 -invariant subspaces of X , and let $\mathcal{N}$ be the subset of Σ consisting of those subspaces which are minimal with respect to inclusion. We note that the member of $\mathcal{N}$ are disjoint and each member of $\mathcal{N}$ is the closed convex hull $C(\Gamma_1 x)$ of some Γ_1 -orbit.

We mainly check corresponding claims in [3, pp.240–241]. From Zorn's lemma, we first obtain Claim 1: $\mathcal{N}$ is non-empty.

For $C_1, C_2 \in \mathcal{N}$, let $p_i : X \rightarrow C_i$ denote the projection of X onto C_i and let $d = d(C_1, C_2) := \inf\{d(x_1, x_2) \mid x_1 \in C_1, x_2 \in C_2\}$. By [3, p.240], we also obtain Claim 2: There exists a unique isometry j of $C_1 \times [0, d]$ onto the convex hull of $C_1 \cup C_2$ such that $j(x, 0) = x$ and $j(x, d) = p_2(x)$ for each $x \in C_1$.

Let $X' = \bigcup \mathcal{N}$. Then we check Claim 3: X' is a closed, convex, Γ -invariant, quasi-dense subspace of X . We note that for each $x \in X'$ there exists a unique member $C \in \mathcal{N}$ such that $x \in C$ (indeed C is the convex hull of the Γ_1 -orbit $\Gamma_1 x$). By [3, p.240], we see that X' is a closed, convex, Γ -invariant subspace of X . We show that X' is quasi-dense in X . Since Γ acts cocompactly on X , $\Gamma B(x_0, N) = X$ for some $x_0 \in X'$ and $N > 0$. Here $\Gamma x_0 \subset X'$ because X' is Γ -invariant. Hence

$$B(X', N) \supset B(\Gamma x_0, N) = \Gamma B(x_0, N) = X.$$

Thus X' is a quasi-dense subspace of X .

Next we check Claim 4: If $C_1, C_2, C_3 \in \mathcal{N}$ and if $p_i : X \rightarrow C_i$ denotes the projection onto C_i , then $p_1 = p_1 p_3$ on C_2 .

Let $f = p_1 p_3 p_2 : C_1 \rightarrow C_2 \rightarrow C_3 \rightarrow C_1$. Then f is an isometry of C_1 by Claim 2. To show that $p_1 = p_1 p_3$ on C_2 , we prove that f is the identity.

We first show that f is an elliptic isometry. Let $K = d(C_1, C_2) + d(C_2, C_3) + d(C_3, C_1)$. Then $d(x, f(x)) \leq K$ for any $x \in C_1$. Hence for each geodesic ray ξ in C_1 , $d(\xi(r), f\xi(r)) \leq K$ for any $r \in [0, \infty)$. This means that every geodesic ray ξ in C_1 is asymptotic to the geodesic ray $f\xi$. Hence the induced homeomorphism $f : \partial C_1 \rightarrow \partial C_1$ is the identity. Thus f is not parabolic isometry by Lemma 2.3. Now we suppose that f is a hyperbolic isometry of C_1 . Then there exists an axis σ_1 of f in C_1 . Let $\sigma_2 = p_2\sigma_1$ and let $\sigma_3 = p_3\sigma_2$. Here we note that σ_2 and σ_3 are geodesic lines in C_2 and C_3 respectively, and the geodesic lines σ_1, σ_2 and σ_3 are parallel by Claim 2. Then

$$p_1\sigma_3 = p_1p_3\sigma_2 = p_1p_3p_2\sigma_1 = f\sigma_1,$$

and $f(\text{Im } \sigma_1) = \text{Im } \sigma_1$. By [3, Lemma II.2.15], f is the identity on $\text{Im } \sigma_1$. This is a contradiction, since σ_1 is an axis of f . Hence f is not a hyperbolic isometry. Thus f is elliptic.

We show that f is the identity. Let $C'_1 = \text{Min}(f)$. Here C'_1 is the fixed point set of f . To show that f is the identity, we prove that $C'_1 = C_1$. If C'_1 is a closed, convex, non-empty, Γ_1 -invariant subspace of X , then $C'_1 = C_1$ by the minimality of C_1 . Now C'_1 is a closed, convex, non-empty subset. Hence it is sufficient to show that C'_1 is Γ_1 -invariant. Let $\gamma_1 \in \Gamma_1$ and let $x \in C'_1$. Then $f(x) = x$. We show that $\gamma_1 x \in C'_1$ (i.e. $f(\gamma_1 x) = \gamma_1 x$). Since p_i is Γ_1 -invariant for each $i = 1, 2, 3$,

$$\begin{aligned} f(\gamma_1 x) &= p_1 p_3 p_2(\gamma_1 x) = p_1 p_3(\gamma_1 p_2(x)) = p_1(\gamma_1 p_3 p_2(x)) \\ &= \gamma_1(p_1 p_3 p_2(x)) = \gamma_1 f(x) = \gamma_1 x. \end{aligned}$$

This means that $\gamma_1 x \in C'_1$. Hence C'_1 is Γ_1 -invariant and $C'_1 = C_1$, i.e., f is the identity.

We fix $X_1 \in \mathcal{N}$ and let $p : X' \rightarrow X_1$ be the orthogonal projection. Let X_2 denote the metric space $(\mathcal{N}, d)$, where $d(C, C') = \inf\{d(x, x') \mid x \in C, x' \in C'\}$. For each $x \in X'$, there exists a unique member $C_x \in \mathcal{N}$ such that $x \in C_x$. Then by [3, p.241], we obtain Claim 5: The map $X' \rightarrow X_1 \times X_2$ given by $x \mapsto (p(x), C_x)$ is a Γ -equivariant isometry.

Here X_1 is Γ_1 -invariant and Γ_1 acts geometrically on X_1 . Finally we show that Γ_2 acts geometrically on X_2 by $\gamma_2 C_x = C_{\gamma_2 x}$ (i.e., Γ_2 acts geometrically on X_2 by projection). It is obvious that the action of Γ_2 on X_2 is cocompact and isometry. Suppose that the action of Γ_2 on X_2 is not proper. Then there exists a number $K > 0$ such that

$$\begin{aligned} &\{\gamma \in \Gamma_2 \mid B(C_{x_0}, K) \cap \gamma B(C_{x_0}, K) \neq \emptyset\} \\ &= \{\gamma \in \Gamma_2 \mid B(C_{x_0}, K) \cap B(C_{\gamma x_0}, K) \neq \emptyset\} \end{aligned}$$

is infinite. Let $\{\gamma_i \mid i \in \mathbb{N}\} = \{\gamma \in \Gamma_2 \mid B(C_{x_0}, K) \cap B(C_{\gamma x_0}, K) \neq \emptyset\}$, where $\gamma_i \neq \gamma_j$ if $i \neq j$. Since $d(C_{x_0}, C_{\gamma_i x_0}) \leq 2K$, there exists a point $y_0 \in C_{x_0}$ such that $d(y_0, \gamma_i x_0) \leq 2K$ by Claim 2. Now Γ_1 acts cocompactly on C_{x_0} (which is the closed convex hull of the orbit $\Gamma_1 x_0$), and $C_{x_0} \subset \Gamma_1 B(x_0, N)$ for some $N > 0$. There exists $\delta_i \in \Gamma_1$ such that $d(\delta_i x_0, y_0) \leq N$. Then

$$d(\delta_i x_0, \gamma_i x_0) \leq d(\delta_i x_0, y_0) + d(y_0, \gamma_i x_0) \leq N + 2K.$$

Thus $d(x_0, \delta_i^{-1} \gamma_i x_0) \leq N + 2K$ for any $i \in \mathbb{N}$. Here we note that $\delta_i^{-1} \gamma_i \neq \delta_j^{-1} \gamma_j$ if $i \neq j$, because $\delta_i, \delta_j \in \Gamma_1, \gamma_i, \gamma_j \in \Gamma_2$ and $\gamma_i \neq \gamma_j$. This means that the action of $\Gamma = \Gamma_1 \times \Gamma_2$ on X is not proper, which is a contradiction. Thus the action of Γ_2 on X_2 is proper, hence, geometric. $\square$

Remark 3.2 We suppose that a group $\Gamma = \Gamma_1 \times \Gamma_2$ acts geometrically on a CAT(0) space X and we suppose that Γ_2 has finite center. Then [3, Lemma II.6.24] implies that Γ_1 acts

cocompactly on the convex hull $C(\Gamma_1 x_0)$ of some Γ_1 -orbit, that is, the condition in Theorem 3.1 holds. In this paper, we mainly use Theorem 3.1 under the condition that Γ_2 has finite center.

From Theorem 3.1 and Lemma 2.4, we obtain the following splitting theorem.

Theorem 3.3 *Suppose that a group $\Gamma = \Gamma_1 \times \Gamma_2$ acts geometrically on a CAT(0) space X . If Γ has finite center, then there exists a closed, convex, Γ -invariant, quasi-dense subspace $X' \subset X$ such that X' splits as a product $X_1 \times X_2$ and the action of $\Gamma = \Gamma_1 \times \Gamma_2$ on $X' = X_1 \times X_2$ is the product action.*

Proof Suppose that Γ has finite center. Then the centers of Γ_1 and Γ_2 are finite. Let X' be a minimal closed convex non-empty Γ -invariant subspace of X (there exists such X' by Zorn's lemma). Then for each $x \in X'$, the convex hull $C(\Gamma x)$ of the Γ -orbit Γx is X' by the minimality of X' . Let $x_0 \in X'$ and let $X_1 = C(\Gamma_1 x_0)$. Since the action of Γ_1 on X_1 is cocompact by [3, Lemma II.6.24], X' splits as $X_1 \times X_2$ by Theorem 3.1. Then we show that $X_2 = C(\Gamma_2 x_0)$ and the action of Γ_2 on X_1 is trivial.

Let $p: X' \rightarrow X_1$ be the projection. For each $\gamma \in \Gamma_2$ and $x \in X_1$, we define $\gamma * x = p(\gamma x)$. Then “ $*$ ” is an isometry action of Γ_2 on X_1 . Here for each $\gamma_1 \in \Gamma_1$, $\gamma_2 \in \Gamma_2$ and $x \in X_1$,

$$\begin{aligned}\gamma_1(\gamma_2 * x) &= \gamma_1 p(\gamma_2 x) = p(\gamma_1 \gamma_2 x) \\ &= p(\gamma_2 \gamma_1 x) = \gamma_2 * (\gamma_1 x).\end{aligned}$$

Hence $\gamma_1(\gamma_2 * x) = \gamma_2 * (\gamma_1 x)$.

Now we show that the orbit $\Gamma_2 * x_0$ is a bounded set. Suppose that $\Gamma_2 * x_0$ is unbounded. Then there exists a sequence $\{g_i * x_0\} \subset \Gamma_2 * x_0$ which converges to some point $\alpha \in \partial X_1$ in $X_1 \cup \partial X_1$. For each $\delta \in \Gamma_1$, $\{\delta(g_i * x_0)\} = \{g_i * (\delta x_0)\}$ converges to α . On the other hand, $\{\delta(g_i * x_0)\}$ converges to $\delta\alpha$. Hence $\delta\alpha = \alpha$ for any $\delta \in \Gamma_1$. This contradicts Lemma 2.4, since the center of Γ_1 is finite. Thus $\Gamma_2 * x_0$ is a bounded set in X_1 . By [3, Corollary II.2.8(1)], there exists a point $y_0 \in X_1$ such that $\Gamma_2 * y_0 = y_0$. Here we may retake the basepoint x_0 as y_0 . Then $X_1 = C(\Gamma_1 x_0)$ and $X_2 = C(\Gamma_2 x_0)$. By the construction of $X_1 \times X_2$ in the proof of Theorem 3.1, the action of Γ_2 on X_1 is trivial. $\square$

A metric space Y is said to be of *non-positive curvature*, if Y is a locally CAT(0) space (cf. [3, p.159]). Let Y be a compact geodesic space of non-positive curvature. Then the universal covering X of Y is a CAT(0) space by the Cartan–Hadamard theorem, and we can think of Y as the quotient $\Gamma \backslash X$ of X , where Γ is the fundamental group of Y acting freely and properly by isometries on X (cf. [3, p. 237]).

As an application of Theorem 3.3, we can obtain the following splitting theorem for compact geodesic spaces of non-positive curvature (i.e. locally CAT(0) spaces). This splitting theorem is relate to the splitting theorems in [3, Theorem II.6.22], [7, 9] and [12].

Theorem 3.4 *Let Y be a compact geodesic space of non-positive curvature (i.e. locally CAT(0)). Suppose that the fundamental group of Y splits as a product $\Gamma = \Gamma_1 \times \Gamma_2$ and that Γ has trivial center. Then there exists a deformation retract Y' of Y which splits as a product $Y_1 \times Y_2$ such that the fundamental group of Y_i is Γ_i for each $i = 1, 2$.*

Proof Let X be the universal covering of Y . Then X is a CAT(0) space, Γ acts freely and properly by isometries on X , and $\Gamma \backslash X = Y$. Since $\Gamma = \Gamma_1 \times \Gamma_2$ acts geometrically on X and Γ has finite center, by Theorem 3.3, there exists a closed, convex, Γ -invariant, quasi-dense

subspace $X' \subset X$ such that X' splits as a product $X_1 \times X_2$ and the action of $\Gamma = \Gamma_1 \times \Gamma_2$ on $X' = X_1 \times X_2$ is the product action. Let $Y_i = \Gamma_i \backslash X_i$ for each $i = 1, 2$. Then

$$\begin{aligned} Y &= \Gamma \backslash X \supset \Gamma \backslash X' = (\Gamma_1 \times \Gamma_2) \backslash (X_1 \times X_2) \\ &= \Gamma_1 \backslash X_1 \times \Gamma_2 \backslash X_2 = Y_1 \times Y_2, \end{aligned}$$

and $\pi_1 Y_i = \Gamma_i$ for each $i = 1, 2$, since Γ_i acts freely and properly by isometries on X_i .

We show that $Y_1 \times Y_2$ is a deformation retract of Y . Since $X' = X_1 \times X_2$ is a quasi-dense subspace of X , there exists $N \geq 0$ such that $B(X', N) = X$. For each $t \in [0, N]$, let $p_t : X \rightarrow B(X', t)$ be the orthogonal projection. Then the map $F : X \times [0, N] \rightarrow X$ defined by $F(x, t) = p_{N-t}(x)$ is a deformation retraction of X' onto X . Since X' is Γ -invariant, $B(X', t)$ is also Γ -invariant and the square

$$\begin{array}{ccc} X & \xrightarrow{p_t} & B(X', t) \\ \gamma \downarrow & & \gamma \downarrow \\ X & \xrightarrow{p_t} & B(X', t) \end{array}$$

is commutative for each $t \in [0, N]$ and $\gamma \in \Gamma$. Hence F induces a deformation retract

$$F' : \Gamma \backslash X \times [0, N] \rightarrow \Gamma \backslash X$$

of $\Gamma \backslash X' = Y_1 \times Y_2$ onto $\Gamma \backslash X = Y$. □

4 A splitting theorem for CAT(0) spaces on which a product group acts geometrically

In this section, we show a splitting theorem in the case a CAT(0) group has infinite center.

We first define an essential subgroup of a group.

Definition 4.1 Let Γ be a group and let G be a subgroup of Γ . In this paper, we say that G is an *essential subgroup* of Γ , if G is a finite-index subgroup of Γ which splits as a product $\Gamma' \times A$, where Γ' has finite center and A is isomorphic to $\mathbb{Z}^n$ for some $n \geq 0$.

The following lemma is known.

Lemma 4.2 ([8, Lemma 2.1]) *If a group $\Gamma_1 \times \Gamma_2$ acts geometrically on some CAT(0) space X , then there exists an essential subgroup of Γ_i for each $i = 1, 2$.*

The purpose of this section is to prove the following main splitting theorem.

Theorem 4.3 *Suppose that a group $\Gamma = \Gamma_1 \times \Gamma_2$ acts geometrically on a CAT(0) space X . Then for essential subgroups G_1 and G_2 of Γ_1 and Γ_2 respectively, there exist closed convex subspaces X_1, X_2, X'_1, X'_2 in X such that*

- (1) $X_1 \times X'_2$ is a closed convex $\Gamma_1 \times G_2$ -invariant quasi-dense subspace of X ,
- (2) $X'_1 \times X_2$ is a closed convex $G_1 \times \Gamma_2$ -invariant quasi-dense subspace of X ,
- (3) X'_1 and X'_2 are isometric to some quasi-dense subspaces of X_1 and X_2 respectively,
- (4) there exist geometric actions of Γ_1 and Γ_2 on X_1 and X_2 respectively, and
- (5) there exist geometric actions of G_1 and G_2 on X'_1 and X'_2 respectively.

Hence the boundary ∂X of X is homeomorphic to the join $\partial X_1 * \partial X_2$ of the boundaries of X_1 and X_2 .

We can obtain the following proposition from Theorems 2.2 and 3.1, Remark 3.2 and Lemma 4.2 by a similar argument to the proof of the splitting theorem in [8]. To prove Theorem 4.3, we investigate details of the argument.

Proposition 4.4 *Suppose that a group $\Gamma = \Gamma_1 \times \Gamma_2$ acts geometrically on a CAT(0) space X . Then for an essential subgroup G_2 of Γ_2 , there exists a closed, convex, $\Gamma_1 \times G_2$ -invariant, quasi-dense subspace X' of X such that X' splits as a product $X_1 \times X'_2$ and there exist geometric actions of Γ_1 and G_2 on X_1 and X'_2 respectively.*

We provide two remarks needed in the proof of Theorem 4.3.

Remark 4.5 Suppose that a group $\Gamma = \Gamma_1 \times \Gamma_2 \times A$ acts geometrically on a CAT(0) space X , where the centers of Γ_1 and Γ_2 are finite and A is isomorphic to $\mathbb{Z}^n$ for some n .

By Theorem 2.2, $\text{Min}(A)$ splits as a product $Z \times Y$ which is a quasi-dense subspace of X , where Z is isometric to $\mathbb{R}^n$, A acts geometrically on Z by restriction and $\Gamma_1 \times \Gamma_2$ acts geometrically on Y by projection. Since $\Gamma_1 \times \Gamma_2$ has finite center, by Theorem 3.3, Y contains a quasi-dense subspace $Y_1 \times Y_2$, where Γ_1 and Γ_2 acts geometrically on Y_1 and Y_2 respectively by the restriction of the action of $\Gamma_1 \times \Gamma_2$ on Y by projection. Here

$$X \supset Z \times Y \supset Z \times Y_1 \times Y_2.$$

We note that for the original action of Γ on X , A acts geometrically on Z by restriction and Γ_i acts geometrically on Y_i by projection for each $i = 1, 2$.

On the other hand, since $\Gamma_1 \times (\Gamma_2 \times A)$ acts geometrically on X and Γ_1 has finite center, by Theorem 3.1 and Remark 3.2, there exists a quasi-dense subspace X' of X which splits as a product $\bar{Y}_1 \times \bar{X}$, where $\Gamma_2 \times A$ acts geometrically on $\bar{X}$ by restriction and Γ_1 acts geometrically on $\bar{Y}_1$ by projection. By Theorem 2.2 (or Theorem 3.1), $\bar{X}$ contains a quasi-dense subspace $\bar{Z} \times \bar{Y}_2$, where A acts geometrically on $\bar{Z}$ by restriction and Γ_2 acts geometrically on $\bar{Y}_2$ by projection. Here

$$X \supset \bar{Y}_1 \times \bar{X} \supset \bar{Y}_1 \times \bar{Z} \times \bar{Y}_2.$$

Also, for the original action of Γ on X , A acts geometrically on Z by restriction and Γ_i acts geometrically on $\bar{Y}_i$ by projection for each $i = 1, 2$.

Then we can identify $Z = \bar{Z}$ because Z and $\bar{Z}$ are the convex hulls of some orbits of A . By construction, $Z \times Y_1 \times Y_2$ and $\bar{Z} \times \bar{Y}_1 \times \bar{Y}_2$ are the unions of all subspaces which are parallel to $Z = \bar{Z}$. Hence we obtain that $Y_1 \times Y_2 = \bar{Y}_1 \times \bar{Y}_2$. Here Γ_i acts geometrically on Y_i and $\bar{Y}_i$ by projection for each $i = 1, 2$. Hence Y_i and $\bar{Y}_i$ are the convex hulls $C(\Gamma_i * x_0)$ of some orbits of Γ_i by the projection action, and we can identify $Y_i = \bar{Y}_i$ for each $i = 1, 2$.

Remark 4.6 Suppose that a group $A_1 \times A_2 \times \Gamma$ acts geometrically on a CAT(0) space X , where A_i is isomorphic to $\mathbb{Z}^{n_i}$ for some n_i ($i = 1, 2$) and Γ has finite center.

By Theorem 2.2, $\text{Min}(A_1 \times A_2)$ is a quasi-dense subspace of X and splits as $Z \times Y$, where $A_1 \times A_2$ acts geometrically on $Z \cong \mathbb{R}^{n_1+n_2}$ by restriction and Γ acts geometrically on Y by projection. Here

$$X \supset \text{Min}(A_1 \times A_2) = Z \times Y.$$

On the other hand, by Theorem 2.2, $\text{Min}(A_1)$ is a quasi-dense subspace of X and splits as $\bar{Z}_1 \times \bar{X}$, where A_1 acts geometrically on $\bar{Z}_1 \cong \mathbb{R}^{n_1}$ by restriction and $A_2 \times \Gamma$ acts geometrically on $\bar{X}$ by projection. Also by Theorem 2.2, $\bar{X}$ contains a quasi-dense subspace $\text{Min}_{\bar{X}}(A_2) = \bar{Z}_2 \times \bar{Y}$, where there exist geometric actions of A_2 and Γ on $\bar{Z}_2 \cong \mathbb{R}^{n_2}$ and $\bar{Y}$,

respectively. Here

$$\begin{aligned} X \supset \text{Min}(A_1) &= \bar{Z}_1 \times \bar{X} \\ &\supset \bar{Z}_1 \times \text{Min}_{\bar{X}}(A_2) = \bar{Z}_1 \times \bar{Z}_2 \times \bar{Y}. \end{aligned}$$

Then by the inductive proof of Theorem 2.2 in [2] and [3, p.245], we can identify $Z = \bar{Z}_1 \times \bar{Z}_2$ and $Y = \bar{Y}$.

Here we note that $Z = \bar{Z}_1 \times \bar{Z}_2$ is the convex hull $C((A_1 \times A_2)x_0)$ of some orbit of $A_1 \times A_2$ and also we note that $Z \times Y = \bar{Z}_1 \times \bar{Z}_2 \times \bar{Y}$ is the union of all subspaces which are parallel to $Z = \bar{Z}_1 \times \bar{Z}_2$ and Γ acts geometrically on $Y = \bar{Y}$ by projection.

Now we prove Theorem 4.3.

Proof of Theorem 4.3 Suppose that a group $\Gamma = \Gamma_1 \times \Gamma_2$ acts geometrically on a CAT(0) space X . Let $G_1 = \Gamma'_1 \times A_1$ and $G_2 = \Gamma'_2 \times A_2$ be essential subgroups of Γ_1 and Γ_2 respectively, where Γ'_i has finite center and A_i is isomorphic to $\mathbb{Z}^{n_i}$ for some n_i for each $i = 1, 2$.

Then $\Gamma'_1 \times A_1 \times \Gamma_2$ is a finite-index subgroup of Γ and acts geometrically on X . Since Γ'_1 has finite center, by Theorem 3.1 and Remark 3.2, X contains a quasi-dense subspace $Y'_1 \times X''_2$, where Γ'_1 acts geometrically on Y'_1 by projection and $A_1 \times \Gamma_2$ acts geometrically on X''_2 by restriction. By Theorem 2.2, $Y'_1 \times X''_2$ contains a quasi-dense subspace

$$Y'_1 \times \text{Min}_{X''_2}(A_1) = \text{Min}_{Y'_1 \times X''_2}(A_1) = Y'_1 \times Z_1 \times X_2,$$

where A_1 acts geometrically on Z_1 by restriction and Γ_2 acts geometrically on X_2 by projection. Here we note that $Y'_1 \times Z_1 \times X_2$ is a closed convex $G_1 \times \Gamma_2$ -invariant quasi-dense subspace of X .

Moreover $G_2 = \Gamma'_2 \times A_2$ is a finite-index subgroup of Γ_2 . Hence $Y'_1 \times Z_1 \times X_2$ contains a quasi-dense subspace $\text{Min}_{Y'_1 \times Z_1 \times X_2}(A_2) = Y'_1 \times Z_1 \times Z_2 \times Y'_2$, where A_2 and Γ'_2 act geometrically on Z_2 and Y'_2 respectively.

Thus

$$\begin{aligned} X \supset Y'_1 \times X''_2 \supset \text{Min}_{Y'_1 \times X''_2}(A_1) &= Y'_1 \times Z_1 \times X_2 \\ &\supset \text{Min}_{Y'_1 \times Z_1 \times X_2}(A_2) = Y'_1 \times Z_1 \times Z_2 \times Y'_2. \end{aligned}$$

Here we note that $Y'_1 \times Z_1 \times Z_2 \times Y'_2$ is a closed convex $G_1 \times G_2$ -invariant quasi-dense subspace of X .

Then we use Remark 4.6. Now $A_1 \times \Gamma_2$ acts geometrically on X''_2 and $A_1 \times A_2 \times \Gamma'_2$ is a finite-index subgroup of $A_1 \times \Gamma_2$. Hence $Y'_1 \times X''_2$ contains a quasi-dense subspace

$$\text{Min}_{Y'_1 \times X''_2}(A_1 \times A_2) = Y'_1 \times \text{Min}_{X''_2}(A_1 \times A_2) = Y'_1 \times \hat{Y}'_2 \times Z,$$

where $A_1 \times A_2$ and Γ'_2 act geometrically on Z and $\hat{Y}'_2$ respectively. Then we can identify $Z = Z_1 \times Z_2$ and $\hat{Y}'_2 = Y'_2$ by Remark 4.6. Thus

$$\begin{aligned} X \supset Y'_1 \times X''_2 \supset \text{Min}_{Y'_1 \times X''_2}(A_1 \times A_2) \\ = Y'_1 \times \text{Min}_{X''_2}(A_1 \times A_2) = Y'_1 \times Y'_2 \times Z_1 \times Z_2. \end{aligned}$$

Next we use Remark 4.5. Since $A_1 \times A_2 \times \Gamma'_1 \times \Gamma'_2$ acts geometrically on X , by Theorem 2.2, X contains a quasi-dense subspace $\text{Min}(A_1 \times A_2) = \bar{Z} \times \bar{Y}'$, where $A_1 \times A_2$ acts geometrically on $\bar{Z}$ by restriction and $\Gamma'_1 \times \Gamma'_2$ acts geometrically on $\bar{Y}'$ by projection. Since $\Gamma'_1 \times \Gamma'_2$ has finite center, by Theorem 3.3, $\bar{Z} \times \bar{Y}'$ contains a quasi-dense subspace

$\bar{Z} \times \bar{Y}'_1 \times \bar{Y}'_2$, where Γ'_i acts geometrically on $\bar{Y}'_i$ for each $i = 1, 2$. Then we can identify $\bar{Z} = Z$, $\bar{Y}'_1 = Y'_1$ and $\bar{Y}'_2 = \hat{Y}'_2$ by Remark 4.5. Thus

$$\begin{aligned} X &\supset \text{Min}(A_1 \times A_2) = \bar{Z} \times \bar{Y}' \\ &\supset \bar{Z} \times \bar{Y}'_1 \times \bar{Y}'_2 = Z \times Y'_1 \times \hat{Y}'_2 \\ &= Z_1 \times Z_2 \times Y'_1 \times Y'_2, \end{aligned}$$

since $Z = Z_1 \times Z_2$ and $\hat{Y}'_2 = Y'_2$. Hence we obtain that

$$X \supset \text{Min}(A_1 \times A_2) \supset Z_1 \times Z_2 \times Y'_1 \times Y'_2. \quad (1)$$

On the other hand, $\Gamma_1 \times A_2 \times \Gamma'_2$ is also a finite-index subgroup of Γ and acts geometrically on X . Since Γ'_2 has finite center, by Theorem 3.1 and Remark 3.2, X contains a quasi-dense subspace $X''_1 \times \tilde{Y}'_2$, where $\Gamma_1 \times A_2$ acts geometrically on X''_1 by restriction and Γ'_2 acts geometrically on $\tilde{Y}'_2$ by projection. By Theorem 2.2, $X''_1 \times \tilde{Y}'_2$ contains a quasi-dense subspace $\text{Min}_{X''_1 \times \tilde{Y}'_2}(A_2) = X_1 \times \tilde{Z}_2 \times \tilde{Y}'_2$, where Γ_1 and A_2 act geometrically on X_1 and $\tilde{Z}_2$ respectively. Here we note that $X_1 \times \tilde{Z}_2 \times \tilde{Y}'_2$ is a closed convex $\Gamma_1 \times G_2$ -invariant quasi-dense subspace of X .

Moreover $G_1 = \Gamma'_1 \times A_1$ is a finite-index subgroup of Γ_1 . Hence $X_1 \times \tilde{Z}_2 \times \tilde{Y}'_2$ contains a quasi-dense subspace $\text{Min}_{X_1 \times \tilde{Z}_2 \times \tilde{Y}'_2}(A_1) = \tilde{Y}'_1 \times \tilde{Z}_1 \times \tilde{Z}_2 \times \tilde{Y}'_2$. Thus

$$\begin{aligned} X &\supset X''_1 \times \tilde{Y}'_2 \supset \text{Min}_{X''_1 \times \tilde{Y}'_2}(A_2) = X_1 \times \tilde{Z}_2 \times \tilde{Y}'_2 \\ &\supset \text{Min}_{X_1 \times \tilde{Z}_2 \times \tilde{Y}'_2}(A_1) = \tilde{Y}'_1 \times \tilde{Z}_1 \times \tilde{Z}_2 \times \tilde{Y}'_2. \end{aligned}$$

By the same argument as the above one, we obtain that

$$X \supset \text{Min}(A_1 \times A_2) \supset \tilde{Z}_1 \times \tilde{Z}_2 \times \tilde{Y}'_1 \times \tilde{Y}'_2. \quad (2)$$

Since $Z_1 \times Z_2$ in (1) and $\tilde{Z}_1 \times \tilde{Z}_2$ in (2) are the convex hulls of some $A_1 \times A_2$ -orbits in X , we can identify $Z_1 \times Z_2 = \tilde{Z}_1 \times \tilde{Z}_2$. Here Z_i and $\tilde{Z}_i$ are isometric to $\mathbb{R}^{n_i}$ for each $i = 1, 2$ by construction. Also $Z_1 \times Z_2 \times Y'_1 \times Y'_2$ in (1) and $\tilde{Z}_1 \times \tilde{Z}_2 \times \tilde{Y}'_1 \times \tilde{Y}'_2$ in (2) are the unions of all subspaces which are parallel to $Z_1 \times Z_2 = \tilde{Z}_1 \times \tilde{Z}_2$ in X , hence, we can identify $Y'_1 \times Y'_2 = \tilde{Y}'_1 \times \tilde{Y}'_2$. Moreover we obtain that $Y'_i = \tilde{Y}'_i$ for each $i = 1, 2$, because Y'_i and $\tilde{Y}'_i$ are the convex hulls of the orbits of the action of Γ'_i on $Y'_1 \times Y'_2 = \tilde{Y}'_1 \times \tilde{Y}'_2$ by projection.

Let $X'_1 = Y'_1 \times Z_1$ and $X'_2 = \tilde{Y}'_2 \times \tilde{Z}_2$. Then

$$X'_1 \times X'_2 = Y'_1 \times Z_1 \times X'_2$$

is a closed convex $G_1 \times G_2$ -invariant quasi-dense subspace of X , and

$$X_1 \times X'_2 = X_1 \times \tilde{Z}_2 \times \tilde{Y}'_2$$

is a closed convex $\Gamma_1 \times G_2$ -invariant quasi-dense subspace of X . Also $X'_1 = Y'_1 \times Z_1$ is isometric to $\tilde{Y}'_1 \times \tilde{Z}_1$ which is a quasi-dense subspace of X_1 , and $X'_2 = \tilde{Y}'_2 \times \tilde{Z}_2$ is isometric to $Y'_2 \times Z_2$ which is a quasi-dense subspace of X_2 . By construction, Γ_i acts geometrically on X_i by projection for each $i = 1, 2$. Also we can obtain a geometric action of $G_i = \Gamma'_i \times A_i$ on X'_i by the product of the geometric action of Γ'_i on $Y'_i = \tilde{Y}'_i$ and the geometric action of A_i on $Z_i \cong \tilde{Z}_i \cong \mathbb{R}^{n_i}$ for each $i = 1, 2$.

Here we obtain that the boundary ∂X_i is homeomorphic to $\partial X'_i$ for each $i = 1, 2$, and

$$\partial X = \partial(X'_1 \times X_2) = \partial X'_1 * \partial X_2 \approx \partial X_1 * \partial X_2.$$

Therefore the boundary ∂X of X is homeomorphic to the join $\partial X_1 * \partial X_2$ of the boundaries of X_1 and X_2 . $\square$

A CAT(0) space X is said to have the *geodesic extension property*, if every geodesic can be extended to a geodesic line $\mathbb{R} \rightarrow X$.

We obtain the following splitting theorem for CAT(0) spaces with the geodesic extension property from [3, Lemma II.6.20] and the argument in the proof of Theorem 4.3.

Theorem 4.7 *Suppose that a group $\Gamma = \Gamma_1 \times \Gamma_2$ acts geometrically on a CAT(0) space X with the geodesic extension property. Then X splits as a product $X_1 \times X_2$ and there exist geometric actions of Γ_1 and Γ_2 on X_1 and X_2 , respectively. Moreover if Γ has finite center, then the action of $\Gamma = \Gamma_1 \times \Gamma_2$ on $X = X_1 \times X_2$ is the product action.*

Proof Suppose that a group $\Gamma = \Gamma_1 \times \Gamma_2$ acts geometrically on a CAT(0) space X with the geodesic extension property. Then by [3, Lemma II.6.20], in the proof of Theorem 4.3, we obtain that $X = X'_1 \times X_2$ where $X'_1 = Y'_1 \times Z_1$, and $X = X_1 \times X'_2 = \tilde{Y}'_1 \times \tilde{Z}_1 \times X'_2$ where $X'_2 = \tilde{Y}'_2 \times \tilde{Z}_2$ and $X_1 = \tilde{Y}'_1 \times \tilde{Z}_1$. Here we note that $X_1 = \tilde{Y}'_1 \times \tilde{Z}_1$ is isometric to $X'_1 = Y'_1 \times Z_1$. Hence there exists a geometric action of Γ_1 on X'_1 . Therefore X splits as a product $X'_1 \times X_2$ and there exist geometric actions of Γ_1 and Γ_2 on X'_1 and X_2 , respectively.

If Γ has finite center, then we obtain that the action splits from Theorem 3.3 and [3, Lemma II.6.20]. $\square$

5 On products of CAT(0) groups

A group Γ is called a *CAT(0) group*, if Γ acts geometrically on some CAT(0) space X . A CAT(0) group Γ is said to be *rigid*, if Γ determines the boundary up to homeomorphisms of a CAT(0) space on which Γ acts geometrically. It is known that there exists a non-rigid CAT(0) group ([5]).

On products of CAT(0) groups, we obtain the following theorem from Theorem 4.3.

Theorem 5.1 *Γ_1 and Γ_2 are CAT(0) groups if and only if $\Gamma_1 \times \Gamma_2$ is a CAT(0) group.*

Proof It is well known that if Γ_1 and Γ_2 are CAT(0) groups then $\Gamma_1 \times \Gamma_2$ is also a CAT(0) group. Indeed if Γ_1 and Γ_2 act geometrically on some CAT(0) spaces X_1 and X_2 respectively, then we can obtain the product geometric action of $\Gamma_1 \times \Gamma_2$ on the CAT(0) space $X_1 \times X_2$.

Conversely, if $\Gamma_1 \times \Gamma_2$ is a CAT(0) group, then $\Gamma_1 \times \Gamma_2$ acts geometrically on some CAT(0) space X and by Theorem 4.3 (4), we obtain that Γ_1 and Γ_2 are CAT(0) groups, because X_1 and X_2 in Theorem 4.3 are CAT(0) spaces. $\square$

On rigidity of products of rigid CAT(0) groups, we also obtain the following theorem from Theorem 4.3.

Theorem 5.2 *If Γ_1 and Γ_2 are rigid CAT(0) groups, then so is $\Gamma_1 \times \Gamma_2$, and the boundary $\partial(\Gamma_1 \times \Gamma_2)$ is homeomorphic to the join $\partial\Gamma_1 * \partial\Gamma_2$ of the boundaries of Γ_1 and Γ_2 .*

Proof Let Γ_1 and Γ_2 be rigid CAT(0) groups. Suppose that $\Gamma_1 \times \Gamma_2$ acts geometrically on a CAT(0) space X . Then by Theorem 4.3, there exist closed convex subspaces X_1 and X_2 in X such that the boundary ∂X is homeomorphic to the join $\partial X_1 * \partial X_2$ and Γ_1 and Γ_2 act geometrically on X_1 and X_2 respectively. Thus $\Gamma_1 \times \Gamma_2$ is a rigid CAT(0) group whose boundary is homeomorphic to the join $\partial\Gamma_1 * \partial\Gamma_2$ of the boundaries of Γ_1 and Γ_2 . $\square$

Open Access This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

References

1. Adams, S., Ballmann, W.: Amenable isometry groups of Hadamard spaces. *Math. Ann.* **312**, 183–195 (1998)
2. Bowers, P., Ruane, K.: Boundaries of nonpositively curved groups of the form $G \times \mathbb{Z}^n$. *Glasgow Math. J.* **38**, 177–189 (1996)
3. Bridson, M.R., Haefliger, A.: *Metric spaces of non-positive curvature*. Springer, Berlin (1999)
4. Burger, M., Schroeder, V.: Amenable groups and stabilizers of measures on the boundary of a Hadamard manifold. *Math. Ann.* **276**, 505–514 (1987)
5. Croke, C.B., Kleiner, B.: Spaces with nonpositive curvature and their ideal boundaries. *Topology* **39**, 549–556 (2000)
6. Ghys, E., de la Harpe, P.: (eds.) *Sur les Groupes Hyperboliques d’apres Mikhael Gromov*. *Progr. Math.* vol. 83, Birkhäuser, Boston (1990).
7. Gromoll, D., Wolf, J.: Some relations between the metric structure and the algebraic structure of the fundamental group in manifolds of non-positive curvature. *Bull. Am. Math. Soc.* **77**, 545–552 (1971)
8. Hosaka, T.: A splitting theorem for CAT(0) spaces with the geodesic extension property. *Tsukuba J. Math.* **27**, 289–293 (2003)
9. Lawson, H.B., Yau, S.T.: Compact manifolds of nonpositive curvature. *J. Differ. Geom.* **7**, 211–228 (1972)
10. Monod, N.: Superrigidity for irreducible lattices and geometric splitting. *J. Am. Math. Soc.* **19**, 781–814 (2006)
11. Ruane, K.: Boundaries of CAT(0) groups of the form $\Gamma = G \times H$. *Topol. Appl.* **92**, 131–152 (1999)
12. Schroeder, V.: A splitting theorem for spaces of nonpositive curvature. *Invent. Math.* **79**, 323–327 (1985)