Erratum to: Abstract supplement Deutsche Gesellschaft für Experimentelle und Klinische Pharmakologie und Toxikologie e.V. Abstracts of the 81th Annual Meeting, March 10-12, 2015, Kiel, Germany

The Publisher¹

Published online: 6 May 2015
© Springer-Verlag Berlin Heidelberg 2015

Erratum to: Naunyn-Schmiedeberg's Arch Pharmacol (2015) 388 (Suppl 1):S1–S98
DOI 10.1007/s00210-015-1087-4

The original version of the abstract 270 on S68 unfortunately contained mistakes. The correct version is given below.

270

Cytotoxicity and Embryotoxicity of a Chinese Herbal Medicine Extraction Used for Atopic Dermatitis

L Li^{1,2,3,4}, A Haase², L Cheng³, CBS Lau³, LY Tang⁴, B Liang⁴, CC Wang⁴, PC Leung³, A Luch², H. Spielmann¹, M Schäfer-Korting¹

¹Institut für Pharmazie, Freie Universität, Berlin, Germany

²Department of Product Safety, German Federal Institute for Risk Assessment (BfR), Berlin, Germany

³Institute of Chinese Medicine, State Key Laboratory of Phytochemistry and Plant Resources in West China, The Chinese University of Hong Kong, Hong Kong,

⁴Department of Obstetrics and Gynaecology, The Chinese University of Hong Kong, Hong Kong
lu.li@fu-berlin.de

Background: Atopic dermatitis (AD) There is considerable interest in traditional Chinese herbal medicines (CHM) as an alternative treatment for AD. A palatable and well tolerated Chinese medicine formula (PHF), composed of five herbal extracts, appears effective in reducing the need for glucocorticoids in moderate-to-severe ADⁱ. No obvious adverse effects were recorded in the clinical studiesⁱⁱ. Yet, further evidence on the safety is still necessary, in particular for the use in pregnancy.

Objectives: The ECVAM validated Embryonic Stem Cell Test (EST)ⁱⁱⁱ is employed to determine the embryotoxicity potential of the PHF formula and the individual components.

Methods & Results: Chromatography technologies (TLC and/or HPLC) were used for chemical authentications to ensure the quality of each test herbal extract. The commercially available permanent mouse embryonic stem cell line D3 and mouse fibroblast cell line 3T3 were applied to the cytotoxicity and differentiation experiments. The embryotoxic potential was derived from the biological and molecular endpoints (ID_{50D3}, IC_{503T3} and IC_{50D3}). According to the linear discriminant functions, both the formula PHF and its individual herbal extracts are classified as non-embryotoxic.

Conclusions: Our study provides evidence on the safety of the formula PHF, which could be used as an adjunctive treatment for AD during pregnancy.

The online version of the original article can be found at :<http://dx.doi.org/10.1007/s00210-015-1087-4>.

✉ The Publisher
content.springerlink@springer.com

¹ Heidelberg, Germany

Acknowledgement:

Deutscher Akademischer Austausch Dienst (DAAD 57076385), Hong Kong General Research Fund (GRF 476210) and National Natural Science Foundation of China (NSFC 81303302).

References:

ⁱHon KL, et al. Therapeutic effect and safety of a traditional Chinese medicine for atopic dermatitis in children: a randomised,

double-blind, placebo-controlled study. *Hong Kong Med J*. 2011;17:S38–40.

ⁱⁱHon KL, et al. A Pentaherbs Capsule as a Treatment Option for Atopic Dermatitis in Children: An Open-Labeled Case Series. *Am J Chin Med*. 2004;32:941–950.

ⁱⁱⁱSeiler AEM and Spielmann H. The validated embryonic stem cell test to predict embryotoxicity in vitro. *Nat Protocols* 2011;6:961–978.