Topiramate raises anterior cingulate cortex glutamine levels in healthy men; a 4.0 T magnetic resonance spectroscopy study

Constance M. Moore · Megan Wardrop ·
Blaise deB. Frederick · Perry F. Renshaw

Published online: 13 December 2006
© Springer-Verlag 2006

Psychopharmacology (2006) 188:236–243

The presentation of Fig. 1b contained errors. The correct figure is given here.

In the Acknowledgements section, the first of the fund numbers (National Institutes of Mental Health) is incorrect. It should be MH01978.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00213-006-0451-y>.

C. M. Moore · M. Wardrop · B. d. Frederick · P. F. Renshaw
Brain Imaging Center, McLean Hospital,
Belmont, MA 02478, USA

C. M. Moore (✉) · B. d. Frederick · P. F. Renshaw
Consolidated Department of Psychiatry, Harvard Medical School,
Boston, MA 02114, USA
e-mail: const@mclean.harvard.edu

Fig. 1 b Proton spectrum (*top*) from the ACC of a 28-year-old male subject 6 h after ingesting 100 mg of topiramate. *Below*, in the LCModel fit (*red line*) and the glutamate (*blue line*), glutamine (*green line*), and GABA (*orange line*) contributions to that LCModel fit a