Erratum to: Comparison of extraction methods for the analysis of natural dyes in historical textiles by high-performance liquid chromatography

Lemonia Valianou · Ioannis Karapanagiotis ·
Yannis Chryssoulakis

Published online: 24 February 2010
© Springer-Verlag 2010

Erratum to: Anal Bioanal Chem DOI 10.1007/s00216-009-3137-6

There was a mistake in Table 1 concerning the absorbance maxima for Rubiadin. The correct values for

Rubiadin are: 243, 277, 411 nm. Also, we note that the Absorbance maxima shown in Table 1 are given in nm. The corrected Table is provided below. Our apologies to the readers.

Table 1 Dyes and colouring compounds investigated in the study. UV-Vis absorbance maxima were measured by HPLC-DAD according to the Experimental

Dye	Dye class [2,36]	Biological source	Main components	Absorbance maxima (nm)
Madder	Mordant dye	<i>Rubia tinctorum</i> L.	alizarin	247, 277, 429
			puruprin	255, 293, 479
			rubiadin	243, 277, 411
Cochineal	Mordant dye	<i>Dactylopius coccus</i> Costa	carminic acid	275, 309, 493
Weld	Mordant dye	<i>Reseda luteola</i> L.	luteolin	253, 265, 345
			luteolin-7-O-glucoside	253, 265, 345
			apigenin	267, 335
Young fustic	Mordant dye	<i>Cotinus coggygria</i> Scop.	fisetin	245, 321, 359
			sulfuretin	257, 267, 377, 397
Turmeric	Direct dye	<i>Curcuma longa</i> L.	curcumin	263, 425
			demethoxycurcumin	249, 421
			bisdemethoxycurcumin	247, 417
Woad	Vat dye	<i>Isatis tinctoria</i> L.	indigotin	241, 285, 330, 605
			indirubin	239, 289, 363, 539

The online version of the original article can be found under <http://dx.doi.org/10.1007/s00216-009-3137-6>.

L. Valianou · I. Karapanagiotis (✉)
Ormylia Art Diagnosis Center,
Ormylia 63071 Chalkidiki, Greece
e-mail: g.karapanagiotis@artdiagnosis.gr

L. Valianou · Y. Chryssoulakis
School of Chemical Engineering,
National Technical University of Athens,
Athens 15773, Greece