Erratum to: Simultaneous determination of NNK and its seven metabolites in rabbit blood by hydrophilic interaction liquid chromatography–tandem mass spectrometry

Hailei Lang · Sheng Wang · Qidong Zhang ·
Beibei Zhao · Lei Wang · Baojun Cao · Juan Wang ·
Jian Mao · Jianxun Zhang

Received: 2 April 2013 / Accepted: 2 April 2013 / Published online: 19 May 2013
© Springer-Verlag Berlin Heidelberg 2013

Erratum to: Analytical and Bioanalytical Chemistry DOI 10.1007/s00216-012-6641-z

The authors would like to call the reader's attention to the fact that unfortunately there was a mistake in Table 3 of this contribution. Please find the correct Table 3 below:

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00216-012-6641-z>.

H. Lang · S. Wang · Q. Zhang · L. Wang · B. Cao · J. Mao ·
J. Zhang (✉)
Zhengzhou Tobacco Research Institute of CNTC,
Zhengzhou 450001, China
e-mail: jxzh258@163.com

B. Zhao
Quality Supervision and Test Station of the Guangxi Tobacco
Monopoly Administration, Nanning 530022, China

J. Wang
Quality Supervision and Test Station of Hainan Tobacco
Corporation, Haikou 571100, China

Table 3 Recovery and RSD of NNK and its metabolites in rabbits plasma

Analyte	Concentration (ng mL ⁻¹)	Mean recovery (%, n=5)	RSD (%, n=5)
NNK	50	107.0	0.85
	10	116.3	5.46
	2	78.9	1.50
HPB	25	101.7	1.99
	5	112.9	5.92
	1	76.9	1.54
NNAL	50	101.4	1.48
	10	99.8	1.56
	2	97.6	3.15
PBD	25	93.6	0.86
	5	107.3	6.52
	1	106.1	4.03
NNK-N-Oxide	25	109.2	0.54
	5	98.2	4.08
	1	93.6	4.21
NNAL-N-Oxide	25	107.0	4.02
	5	104.4	3.03
	1	102.3	4.23
Keto acid	25	92.0	0.53
	5	101.7	1.99
	1	82.3	1.81
Hydroxy acid	25	109.2	0.54
	5	108.8	5.94
	1	90.8	4.10