Erratum to: Enantioseparation of flinderoles and borreverines by HPLC on Chirobiotic V and V2 stationary phases and by CE using cyclodextrin selectors

Jonathan P. Smuts · Yun-Cheol Na ·
Ravikrishna Vallakati · Adam Přibylka ·
Jeremy A. May · Daniel W. Armstrong

Received: 26 November 2013 / Accepted: 6 December 2013 / Published online: 10 January 2014
© Springer-Verlag Berlin Heidelberg 2014

Erratum to: Analytical and Bioanalytical Chemistry
DOI 10.1007/s00216-013-7335-x

Regrettably there was an error in Fig. 2 where the structures of flinderole B and C have been drawn with the stereogenic methyl group adjacent to the stereogenic isoprenyl group. The correct representation of Fig. 2 is shown below as may also be found by comparing with Fig. 1 of our publication. The conclusion drawn from the figure in the publication, i.e. “in the *anti* configuration both faces of the flinderole are blocked for π - π and hydrogen bonding interactions whereas in the *syn* configuration one face is more accessible”, remains unchanged.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00216-013-7335-x>.

J. P. Smuts · D. W. Armstrong (✉)
Department of Chemistry and Biochemistry, University of Texas at
Arlington, Arlington, TX 76019, USA
e-mail: sec4dwa@uta.edu

Y.-C. Na
Seoul Center, Korea Basic Science Institute, 136701 Seoul,
South Korea

R. Vallakati · J. A. May
Department of Chemistry, University of Houston, 136 Fleming
Building, Houston, TX 77204-5003, USA

A. Přibylka
Regional Centre of Advanced Technologies and Materials,
Department of Analytical Chemistry, Palacký University,
17. Listopadu 12, 77146 Olomouc, Czech Republic

Fig. 2 A comparison of the two- and three-dimensional representations of flinderole B and C and the impact of the *syn* versus *anti* configuration of the methyl and isoprenyl groups on potential chiral recognition. Three-dimensional optimizations were conducted with ACD ChemsSketch freeware