**Franka Kahlenberg · Daniel Sanchez ·
Ingolf Lachmann · Ludmila Tuckova ·
Helena Tlaskalova · Enrique Méndez ·
Thomas Mothes**

Monoclonal antibody R5 for detection of putatively coeliac-toxic gliadin peptides

Published online: 21 February 2006
© Springer-Verlag 2006

**Eur Food Res Technol (2006) 222: 78–82
DOI 10.1007/s00217-005-0100-4**

Due to an unfortunate error, the text on page 80, right-hand column, line 4 was incorrect. The text should read:

“tTG is able to bind biotin-cadaverine to the second amino acid residue in the PQQFP motif but not to other residues (results not shown).”

The online version of the original article can be found at
<http://dx.doi.org/10.1007/s00217-005-0100-4>

F. Kahlenberg · T. Mothes (✉)
Institute of Laboratory Medicine, Clinical Chemistry, and
Molecular Diagnostics, University Hospital Leipzig,
Leipzig, Germany
e-mail: mothes@medizin.uni-leipzig.de
Tel.: +49-341-9722251
Fax: +49-341-9722209

D. Sanchez · L. Tuckova · H. Tlaskalova
Department of Immunology, Institute of Microbiology, Czech
Academy of Sciences,
Prague, Czech Republic

I. Lachmann
Roboscreen GmbH Leipzig,
Leipzig, Germany

E. Méndez
Unidad de Gluten, Centre Nacional de Biotecnologia, CSIC,
Campus Universidad Autonoma,
Cantoblanco, Madrid, Spain