Oligopeptides hydrolysed by muscle dipeptidyl peptidases can generate angiotensin-I converting enzyme inhibitory dipeptides

Miguel Ángel Sentandreu · Fidel Toldrá

Published online: 15 August 2006
© Springer-Verlag 2006

Eur Food Res Technol (2006) s00217-006-0367-0

Unfortunately, Table was published with errors. The correct table is given here:

Table 3 IC₅₀ values of dipeptides generated by the action of muscle DPP on the assayed substrates

Peptide	IC ₅₀ ^a (μM)
Gly-Gly	3450.4 ± 208.4
Ala-Gly	1020.2 ± 85.2
Phe-Gly	515.7 ± 0.30
Arg-Gly	362.2 ± 14.6
Tyr-Gly	330.0 ± 20.6
Val-Gly	149.2 ± 5.7
Met-Ala	152.0 ± 5.2
Arg-Phe	136.7 ± 5.4
Gly-Phe	117.1 ± 2.8
Arg-Ser	61.5 ± 5.9
Val-Tyr	4.6 ± 0.2

Results expressed as the mean of three replicates ± standard deviation.

^aThe concentration inhibiting 50% of angiotensin-I converting enzyme activity, determined as described in materials and methods.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00217-006-0367-0>.

M. Á. Sentandreu (✉) · F. Toldrá
Instituto de Agroquímica y Tecnología de Alimentos (CSIC),
P.O. Box 73, 46100 Burjassot, Valencia, Spain
e-mail: ciesen@iata.csic.es
Fax: + 34-3636301