Arnaud Fumal · Valentin Bohotin ·
Michel Vandenheede · Laurence Seidel ·
Victor De Pasqua · Alain Maertens de Noordhout ·
Jean Schoenen

Effects of repetitive transcranial magnetic stimulation on visual evoked potentials: new insights in healthy subjects

Published online: 12 June 2003
© Springer-Verlag 2003

Exp Brain Res (2003) 150:332–340

In the original article, Fig. 3 was published incorrect. The correct figure is shown below.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00221-003-1423-7>

A. Fumal · V. Bohotin · M. Vandenheede · V. De Pasqua ·
A. M. de Noordhout · J. Schoenen (✉)
University Department of Neurology,
CHR Citadelle Hospital,
4000 Liège, Belgium
e-mail: jschoenen@ulg.ac.be
Tel.: +32-4-2256111
Fax: +32-4-2256451

A. Fumal · J. Schoenen
Department of Neuroanatomy,
University of Liège,
4020 Liège, Belgium

L. Seidel
Department of Medical Informatics, CHU Hospital,
University of Liège,
4000 Liège, Belgium

Fig. 3 Poststimulus changes in first block amplitude (μV, *upper*), habituation (% *middle*) and habituation slope (*lower*) of the N1-P1 component of PR-VEP in three rTMS sessions at 1 or 10 Hz repeated in the same ten subjects (mean ± SD)