Damage to superior parietal cortex impairs pointing in the sagittal Plane

James Danckert · Lana Goldberg · Carol Broderick

Published online: 14 April 2009
© Springer-Verlag 2009

Erratum to: Exp Brain Res DOI 10.1007/s00221-009-1766-9

The author regrets to inform that there is a mistake in Table 1, the value 48.9 for patient ME's mean under the right hand "away" column is incorrect, it should be 56.8.

Table 1 with the corrected value is given below.

Table 1 *T* values (modiWed Crawford method: Crawford and Howell 1998; Crawford and Garthwaite 2002) with associated *P* values and the proportion of the normal population below ME's score (*P* below) for the time spent post-peak velocity

Movement direction	Left hand				Right hand			
	Left	Right	Towards	Away	Left	Right	Towards	Away
Control mean (SD)	48.92 (3.1)	51 (2.74)	51.81 (2.4)	49.6 (3.04)	50.2 (2.8)	48.6 (2.4)	52.2 (1.06)	48.9 (2.6)
ME mean	54	59.075	60.2	60.225	54.425	51.525	57.4	56.8
<i>t</i> value (1-tailed)	1.55 (<i>P</i> = 0.08)	2.78 (<i>P</i> = 0.014)	3.22 (<i>P</i> = 0.007)	3.29 (<i>P</i> = 0.007)	1.42 (<i>P</i> = 0.09)	1.15 (<i>P</i> = 0.14)	2.001 (<i>P</i> = 0.04)	3.23 (<i>P</i> = 0.007)
<i>P</i> below	91.69	98.63	99.26	99.34	90.11	85.59	95.73	99.27

The online version of the original article can be found under doi:[10.1007/s00221-009-1766-9](https://doi.org/10.1007/s00221-009-1766-9).

J. Danckert (✉) · L. Goldberg · C. Broderick
Department of Psychology, University of Waterloo,
200 University Avenue West, Waterloo,
ON N2L 3G1, Canada
e-mail: jdanker@watarts.uwaterloo.ca; jdanker@uwaterloo.ca