Erratum to: Dependency of human neck reflex responses on the bandwidth of pseudorandom anterior-posterior torso perturbations

Patrick A. Forbes · Edo de Bruijn · Alfred C. Schouten ·
Frans C. T. van der Helm · Riender Happee

/ Published online: 23 March 2013
© Springer-Verlag Berlin Heidelberg 2013

Erratum to: Exp Brain Res
DOI 10.1007/s00221-012-3388-x

In the original publication of this article, a labelling error was made in plotting the gain of the $\dot{\theta}_H$ kFRFs in Figs. 4, 6, and 7 (top-middle plot). The label on the y axis was in the units $^{\circ}/s/m/s$; however, the lines were plotted in the units radians/s/m/s. To maintain unit consistency in plots throughout the manuscript, these errors are fixed by reprinting the figures with the label and lines having the same scale ($^{\circ}/s/m/s$).

The revised figures are reproduced below:

The online version of the original article can be found under doi: [10.1007/s00221-012-3388-x](https://doi.org/10.1007/s00221-012-3388-x).

P. A. Forbes (✉) · E. de Bruijn · A. C. Schouten ·
F. C. T. van der Helm · R. Happee
Department of Biomechanical Engineering,
Faculty of Mechanical, Maritime and Materials Engineering,
Delft University of Technology, Mekelweg 2,
2628 CD Delft, The Netherlands
e-mail: p.a.forbes@tudelft.nl

A. C. Schouten · F. C. T. van der Helm
Laboratory of Biomechanical Engineering, Institute
for Biomedical Technology and Technical Medicine (MIRA),
University of Twente, P.O. Box 217, 7500 EA Enschede,
The Netherlands

Fig. 4 Bandwidth variation-related kFRFs of head global (X_{GH}), rotational (θ_H), and relative (X_{RH}) velocity (left-to-right) averaged across all subjects (shaded region is $\pm$ SE) for four bandwidth

conditions at amplitude A2 with eyes open (EO). Gains were log transformed before calculating the subject average and SE

Fig. 6 Amplitude variation-related kFRFs (gain and phase only) of head global (X_{GH}), rotational (θ_H), and relative (X_{RH}) velocity (left-to-right) averaged across all subjects (shaded region is $\pm$ SE) for

amplitudes A1/A2/A3 during perturbation V4 (bandwidth = 4 Hz). Gains were log transformed before calculating the subject average and SE

Fig. 7 EO/EC variation-related kFRFs (gain and phase only) of head global (X_{GH}), rotational (θ_H), and relative (X_{RH}) velocity (left-to-right) averaged across all subjects (shaded region is $\pm$ SE) for amplitude A_2

during perturbation V4 (bandwidth = 4 Hz). Gains were log transformed before calculating the subject average and SE