Arata Horii · Paul F. Smith · Cynthia L. Darlington

Quantitative changes in gene expression of glutamate receptor subunits/subtypes in the vestibular nucleus, inferior olive and flocculus before and following unilateral labyrinthectomy in the rat: real-time quantitative PCR method

Published online: 20 July 2001
© Springer-Verlag 2001

Exp Brain Res (2001)139:188–200

The two footnotes to Table 4 were rendered incorrectly.
They should have been given as follows:

^a $P < 0.05$ vs left

^b $P < 0.05$ vs 50 h postsham

The online version of the original article can be found at
<http://dx.doi.org/10.1007/s002210100763>

A. Horii (✉) · P.F. Smith
Vestibular Research Group, Department of Pharmacology, School
of Medical Sciences, University of Otago Medical School, Dun-
edin, New Zealand

A. Horii
Department of Otolaryngology and Sensory Organ Surgery, Osaka
University Medical School, Osaka, Japan

C.L. Darlington
Vestibular Research Group, Department of Psychology and the
Neuroscience Research Centre, University of Otago, Dunedin,
New Zealand

Present address:

Arata Horii,
Department of Otolaryngology and Sensory Organ Surgery,
Osaka University Medical School, Suita,
Osaka 565-0871, Japan
e-mail: ahorii@ent.med.osaka-u.ac.jp
Tel.: +81-6-68793951, Fax: +81-6-68793959