Erratum

Calcified tissue International Vol. 68, No. 2, February 2001, pp. 109–116 DOI: 10.1007/s002230001191)

Table 4 was published correctly in the electronic version of “Quantification of the Expression Levels of Lysosomal Cysteine Proteinases in Purified Human Osteoclastic Cells by Competitive RT-PCR, by O. Ishibashi, T. Inui, Y. Mori, T. Kurokawa, T. Kokubo, and M. Kumegawa. However, the Table 4 that appeared in the print version was incorrect. It appears correctly below.

Table 4. Expression levels of lysosomal cysteine proteinases in human osteoplast-like cells isolated from giant cell tumor of bone and bone marrow cells in culture^a

Source of OLCs	GCT		BMC	
	Cathepsin/ GAPDH	Relative rate	Cathepsin/ GAPDH	Relative rate
Cathepsin K	14	100	9.6	100
Cathepsin B	0.72	5.1	0.73	7.5
Cathepsin L	0.1	0.76	0.0021	0.022
Cathepsin S	0.034	0.24	0.0069	0.072
GAPDH	(1)	7.1	(1)	10

^a For each of the GCT- and BMC-derived OLCs, the mRNA levels of cathepsins B, K, L, and S are normalized for the GAPDH mRNA level, and results are shown in the left column. The relative levels of the normalized values are shown in the right columns (the values for cathepsin K are set at 100).