Comparisons of genetic population structures in four intertidal brachyuran species of contrasting habitat characteristics

Masako Kawane · Keiji Wada · Katsutoshi Watanabe

Published online: 28 November 2008
© Springer-Verlag 2008

Erratum to: Mar Biol (2008) 156:193–203
DOI 10.1007/s00227-008-1076-y

Unfortunately, the presentation of Tables 1, 2, 3, and 4 contain errors. The correct tables are given below.

The online version of the original article can be found under
doi: [10.1007/s00227-008-1076-y](https://doi.org/10.1007/s00227-008-1076-y).

M. Kawane · K. Wada (✉)
Department of Biological Science, Faculty of Science,
Nara Women's University, Kitauoya-nishimachi,
Nara 630-8506, Japan
e-mail: mbanzai@cc.nara-wu.ac.jp

K. Watanabe
Laboratory of Animal Ecology, Graduate School of Science,
Kyoto University, Kitashirakawa-Oiwakecho,
Sakyo-ku, Kyoto 606-8502, Japan

Table 1 Haplotype composition, haplotype diversity and nucleotide diversity of 10 *Ocypode ceratophthalma* population

Locality number	Haplotype														
	Oc1	Oc2	Oc3	Oc4	Oc5	Oc6	Oc7	Oc8	Oc9	Oc10	Oc11	Oc12	Oc13	Oc14	Oc15
6	4	1	2	1	2										
9	2	1		4	1	1	1								
16	4		2	1	1			1	1						
18	3			2	3					1	1				
23	2			1	1							1	1	1	1
27	1			2	1										
32	3			2	3										
33	2		1	4											
34	1			1											
35	1														
Total	23	2	5	18	12	1	1	1	1	1	1	1	1	1	1

Locality number	Haplotype											<i>n</i>	Haplotype diversity	Nucleotide diversity
	Oc16	Oc17	Oc18	Oc19	Oc20	Oc21	Oc22	Oc23	Oc24	Oc25	Oc26			
6												10	0.822 ± 0.097	0.0022 ± 0.0018
9												10	0.844 ± 0.103	0.0035 ± 0.0025
16												10	0.844 ± 0.103	0.0085 ± 0.0052
18												10	0.844 ± 0.080	0.0033 ± 0.0024
23	1	1										10	0.978 ± 0.054	0.0043 ± 0.0029
27			1									5	0.900 ± 0.161	0.0028 ± 0.0024
32				1	1							10	0.844 ± 0.080	0.0026 ± 0.0020
33	1		1			1						10	0.844 ± 0.103	0.0031 ± 0.0023
34							1	1	1			5	1.000 ± 0.127	0.0053 ± 0.0039
35										1	3	5	0.700 ± 0.218	0.0203 ± 0.0131
Total	2	1	2	1	1	1	1	1	1	1	3	85	0.862 ± 0.084	0.0056 ± 0.0055

Locality numbers as given in Fig. 1

Table 2 Haplotype composition, haplotype diversity and nucleotide diversity of 10 *Gaetice depressus* population

Locality number	Haplotype														
	Gd1	Gd2	Gd3	Gd4	Gd5	Gd6	Gd7	Gd8	Gd9	Gd10	Gd11	Gd12	Gd13	Gd14	Gd15
1	5	1	3	1	1	1									
3	3		3				1	1	1	1					
8	5		2							1	1	1			
14	4		2										1	1	1
18	1		3		1										
22	1		7												
28	4		5												
31															
Total	21	1	25	1	2	1	1	1	1	2	1	1	1	1	1

Locality number	Haplotype												<i>n</i>	Haplotype diversity	Nucleotide diversity
	Gd16	Gd17	Gd18	Gd19	Gd20	Gd21	Gd22	Gd23	Gd24	Gd25	Gd26	Gd27			
1													10	0.867 ± 0.085	0.0027 ± 0.0021
3													10	0.867 ± 0.085	0.0036 ± 0.0025
8													10	0.756 ± 0.130	0.0034 ± 0.0025
14	1												10	0.844 ± 0.103	0.0031 ± 0.0023
18		1	1	1	1	1							10	0.933 ± 0.077	0.0042 ± 0.0029
22							1	1					10	0.533 ± 0.180	0.0012 ± 0.0012
28									1				10	0.644 ± 0.101	0.0015 ± 0.0014
31										7	2	1	10	0.511 ± 0.164	0.0011 ± 0.0011
Total	1	1	1	1	1	1	1	1	1	7	2	1	80	0.744 ± 0.163	0.0026 ± 0.0012

Locality numbers as given in Fig. 1

Table 3 Haplotype composition, haplotype diversity and nucleotide diversity of 10 *Chiromantes dehaani* population

Locality number	Haplotype													
	Cd1	Cd2	Cd3	Cd4	Cd5	Cd6	Cd7	Cd8	Cd9	Cd10	Cd11	Cd12	Cd13	Cd14
2	4	1	1	2	1	1								
4	4		1	1			1	1	1	1				
3	4	1	2			1	1				1			
15	1					1				1		1	1	1
17	3			1								1		
20	4			1		2			1					1
24	4		3	1	1	1								
26	4		2	2										
30	4			2										
Total	32	2	9	10	2	6	2	1	2	2	1	2	1	2

Locality number	Haplotype											<i>n</i>	Haplotype diversity	Nucleotide diversity
	Cd15	Cd16	Cd17	Cd18	Cd19	Cd20	Cd21	Cd22	Cd23	Cd24	Cd25			
2												10	0.844 ± 0.103	0.0055 ± 0.0036
4												10	0.867 ± 0.107	0.0054 ± 0.0036
3												10	0.844 ± 0.103	0.0052 ± 0.0035
15	1	1	1	1								10	1.000 ± 0.045	0.0081 ± 0.0050
17												5	0.700 ± 0.218	0.0016 ± 0.0016
20					1							10	0.844 ± 0.103	0.0061 ± 0.0039
24												10	0.800 ± 0.100	0.0046 ± 0.0031
26						1	1					10	0.822 ± 0.097	0.0042 ± 0.0029
30								1	1	1	1	10	0.844 ± 0.103	0.0041 ± 0.0028
Total	1	1	1	1	1	1	1	1	1	1	1	85	0.841 ± 0.077	0.0050 ± 0.0017

Locality numbers as given in Fig. 1

Table 4 Haplotype composition, haplotype diversity and nucleotide diversity of 10 *Deiratonotus japonicus* population

Locality number	Haplotype																		
	Dj1	Dj2	Dj3	Dj4	Dj5	Dj6	Dj7	Dj8	Dj9	Dj10	Dj11	Dj12	Dj13	Dj14	Dj15	Dj16	Dj17	Dj18	Dj19
5	2	1	2																
14			1	2	1	1													
11			3	2			1	1	1	1	1								
12			1	1								1	6	1	1				
13			4	3									2			1	2	1	1
15																			
21			2																
24			1																
25																			
29																			
Total	2	1	14	8	1	1	1	1	1	1	1	1	8	1	1	1	2	1	1

Locality number	Haplotype															<i>n</i>	Haplotype diversity	Nucleotide diversity
	Dj20	Dj21	Dj22	Dj23	Dj24	Dj25	Dj26	Dj27	Dj28	Dj29	Dj30	Dj31	Dj32	Dj33	Dj34			
5																5	0.800 ± 0.164	0.0019 ± 0.0017
14																5	0.900 ± 0.161	0.0022 ± 0.0020
11																10	0.911 ± 0.077	0.0032 ± 0.0023
12																11	0.727 ± 0.144	0.0035 ± 0.0024
13	1															15	0.895 ± 0.053	0.0038 ± 0.0026
15		1	1	1	1	1										5	1.000 ± 0.127	0.0041 ± 0.0032
21						1	1	1								5	0.900 ± 0.161	0.0064 ± 0.0045
24									1	1	1	1				5	1.000 ± 0.127	0.0048 ± 0.0036
25								1					4			5	0.400 ± 0.237	0.0030 ± 0.0025
29														1	4	5	0.400 ± 0.237	0.0007 ± 0.0009
Total	1	1	1	1	1	2	1	2	1	1	1	1	4	1	4	71	0.793 ± 0.223	0.0034 ± 0.0016

Locality numbers as given in Fig. 1