Asymptotic Nusselt and entropy generation numbers for non-Newtonian fluid flow

Shohel Mahmud · Roydon Andrew Fraser

Received: 23 January 2006 / Accepted: 9 February 2006 / Published online: 26 October 2006
© Springer-Verlag 2006

Heat Mass Transfer (2005) 41(11):999–1003

The authors regret that the following typos occurred in the above article:

1. Equation 7: The right-hand side should be multiplied by 4 instead of the left-hand side

$$U \frac{\partial \Theta}{\partial X} = 4 \frac{1}{R^j} \frac{\partial}{\partial R} \left(R^j \frac{\partial \Theta}{\partial R} \right). \quad (7)$$

2. Equation 9: The second term on the right-hand side should be multiplied by 16

$$\Theta = \frac{(m+j+1)(j+1)}{m} \left[\frac{R^2}{2(j+1)} - \frac{R^{m+2}}{(m+j+1)(m+2)} \right] + \frac{16X}{2(2-j)} + \Gamma. \quad (9)$$

3. Equation 13: The right-hand side should be multiplied by 16

$$\Theta_m = -\frac{16}{2} \frac{X}{j-2}. \quad (13)$$

4. Equation 16: $15j^2$ at the denominator should be replaced by $5j^2$

$$\text{Nu}_\infty = \frac{2(2-j)(3+j)(3+m+j)(3+2m+j)}{2m^2 + 7jm + 13m + 17 + 18j + 5j^2}. \quad (16)$$

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00231-005-0622-0>.

S. Mahmud (✉) · R. A. Fraser
Department of Mechanical Engineering,
University of Waterloo,
Waterloo, ON N2L3G1, Canada
e-mail: smahmud@uwaterloo.ca