Early anisotropy changes in the corpus callosum of patients with optic neuritis

M. Bester · C. Heesen · S. Schippling · R. Martin ·
X.-Q. Ding · B. Holst · J. Fiehler

Published online: 11 September 2008
© Springer-Verlag 2008

Erratum to: Neuroradiology (2008) 50:549–557
DOI 10.1007/s00234-008-0377-7

An erroneous number for the mean FA of the GCC in the control group was reported at MRI3 in the section “Follow up of white matter lesions, ADC and FA”. It should read (0.72 vs 0.76, $P < 0.01$). Due to an oversight wrong ADC and FA values for the control group were reported in Table 2. The table should read as follows:

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00234-008-0377-7>.

M. Bester · C. Heesen · S. Schippling · R. Martin · X.-Q. Ding ·
B. Holst · J. Fiehler (✉)
Department of Neuroradiology,
University Medical Center Hamburg-Eppendorf,
Martinistrasse 52,
20246 Hamburg, Germany
e-mail: fiehler@uke.uni-hamburg.de

C. Heesen · S. Schippling · R. Martin
Institute for Neuroimmunology and Clinical MS Research,
University Medical Center Hamburg-Eppendorf,
Hamburg, Germany

Table 2 ADC ($10^{-3}\text{mm}^2/\text{s}$) and FA values of patients and controls at many specified time

	MRI														
	1			2			3			4			Control		
	Mean	n	SD	Mean	n	SD	Mean	n	SD	Mean	n	SD	Mean	n	SD
FWM ADC	77.61	24	9.03	75.91	24	9.60	78.91	15	7.39	82.66	5	8.00	74.27	15	4.28
Genu ADC	79.88	24	7.52	81.64	24	10.24	81.15	15	7.18	85.52	5	8.09	81.78	15	7.83
OWM ADC	85.60	24	11.44	85.85	24	10.71	84.37	15	8.66	92.03	5	7.24	82.13	15	4.52
Splenium ADC	78.55	24	7.30	77.50	24	5.51	81.60	15	8.60	79.99	5	5.94	77.54	15	7.03
FWM FA	0.448	24	0.11	0.390	24	0.13	0.436	15	0.07	0.394	5	0.12	0.401	15	0.06
Genu FA	0.725	24	0.09	0.714	24	0.10	0.721	15	0.08	0.619	5	0.07	0.764	15	0.06
OWM FA	0.520	24	0.16	0.527	24	0.15	0.502	15	0.16	0.463	5	0.14	0.566	15	0.07
Splenium FA	0.726	24	0.08	0.691	24	0.12	0.715	15	0.09	0.629	5	0.08	0.816	15	0.03