The Use of Feathers to Monitor Heavy Metal Contamination in Herons, Korea

Jungsoo Kim · Tae-Hoe Koo

Published online: 5 December 2007
© Springer Science+Business Media, LLC 2007

Erratum to: Arch Environ Contam Toxicol (2007) 53:435–441
DOI 10.1007/s00244-006-0196-y

In the online and printed version of this article, the mean $\pm$ SD values of the Grey Heron in Table 4 appear incorrectly. They appear correctly below:

Table 4 Concentrations (mean $\pm$ SD, $\mu\text{g/wet g}$) of heavy metals in feathers of Black-crowned Night Heron and Grey Heron chicks

Species	Zinc	Manganese	Iron	Copper	Lead	Cadmium ^a
Black-crowned Night Heron ($n = 8$)						
Frequency	8	8	8	8	8	8
Mean $\pm$ SD	38.3 $\pm$ 5.34	27.0 $\pm$ 29.8	45.8 $\pm$ 24.0	9.93 $\pm$ 2.26	0.33 $\pm$ 0.11	150 $\pm$ 88.2
Min-max	29.8–46.5	3.50–88.2	27.3–95.2	6.76–12.8	0.21–0.52	21.5–324
Geomean	38.0	16.0	41.6	9.70	0.31	121
Grey Heron ($n = 9$)						
Frequency	9	9	9	9	9	9
Mean $\pm$ SD	50.9 $\pm$ 14.0	21.0 $\pm$ 16.3	155 $\pm$ 123	30.2 $\pm$ 12.9	0.25 $\pm$ 0.07	143 $\pm$ 32.0
Min-max	33.2–74.7	4.31–56.8	59.1–420	14.3–49.2	0.17–0.41	83.0–186
Geomean	49.3	16.2	125	27.7	0.24	139
P value ^b	NS	NS	0.001	>0.00001	0.05	NS

^a Was $\mu\text{g/wet kg}$

^b p value for t test comparing between species

NS = not significant

The online version of the original article can be found under doi:
[10.1007/s00244-006-0196-y](https://doi.org/10.1007/s00244-006-0196-y)

J. Kim
Center for Environmental Studies, Kyung Hee University,
446-701 Yongin, Gyeonggi-do, Republic of Korea

T.-H. Koo (✉)
School of Environmental and Applied Chemistry, Kyung Hee
University, 446-701 Yongin, Gyeonggi-do, Republic of Korea
e-mail: thkoo@khu.ac.kr