Heavy Metal Concentrations in Feathers of Korean Shorebirds

Jungsoo Kim · Tae-Hoe Koo

Published online: 4 March 2008
© Springer Science+Business Media, LLC 2008

Erratum to: Arch Environ Contam Toxicol (2008)
DOI 10.1007/s00244-007-9089-y

In the online and printed version of this article, the *p*-values in Table 3 are incorrect. They appear correctly below:

Table 3 Heavy metal concentrations (geomean, 95% CI µg/g wet weight) in the liver of shorebirds from Yeongjong Island, Korea

Species		Zinc	Manganese	Copper	Lead	Cadmium
Kentish Plover (<i>n</i> = 5)	Mean	26.6	1.35	5.51	4.26a	0.55
	Extreme	10.6–42.6	0.01–3.02	1.83–9.20	1.19–7.34	0.35–0.75
Mongolian Plover ^a (<i>n</i> = 2)	Mean	30.9	1.07	5.34	2.05	0.14
	Extreme	23.9–38.0	0.01–3.44	3.58–7.12	ND–2.05	0.01–0.36
Dunlin (<i>n</i> = 6)	Mean	39.5	3.05	8.64	3.64a	0.67
	Extreme	35.4–43.6	2.43–3.66	1.35–15.9	2.78–4.49	0.46–0.88
Great Knot (<i>n</i> = 10)	Mean	31.6	1.11	3.52	3.37a	0.48
	Extreme	27.3–35.9	0.65–1.56	0.01–9.87	1.39–5.35	0.13–0.83
Terek Sandpiper (<i>n</i> = 10)	Mean	20.6	1.61	6.36	0.85b	0.34
	Extreme	13.5–27.8	0.96–2.26	2.19–10.5	0.20–1.51	0.14–0.54
<i>p</i> -Value		NS	NS	NS	0.001	NS

Note: Means sharing same letter were not significantly different among species. ND = not detected, NS = not significant

^a The Mongolian Plover was excluded from statistical analyses because of the small sample size

The online version of the original article can be found under doi:
[10.1007/s00244-007-9089-y](https://doi.org/10.1007/s00244-007-9089-y).

J. Kim · T.-H. Koo (✉)
Department of Environmental Science and Engineering,
Kyung Hee University, Yongin, Gyeonggi-do 446-701,
Republic of Korea
e-mail: thkoo@khu.ac.kr