Erratum to: Prenatal Diagnosis and Management of Fetal Long QT Syndrome

Viktor Tomek · Jan Skovranek · Roman A. Gebauer

Published online: 26 September 2009
© Springer Science+Business Media, LLC 2009

Erratum to: *Pediatr Cardiol* (2009) 30:194–196 DOI 10.1007/s00246-008-9294-0

In the online and print versions of this article published previously, Fig. 2 was presented incorrectly. In the figure of the pulsed Doppler echocardiogram depicting a second degree A-V block, there were interchanged signs of short

early deceleration time of mitral valve flow (arrows) and atrial contraction wave (A) in beats with preserved atrio-ventricular conduction. In the correctly annotated image, short early deceleration time of mitral valve flow (arrows) is followed by atrial contraction wave (A). The corrected figure and caption appear below.

Fig. 2 Pulsed Doppler echocardiogram showing a very short early deceleration time of mitral valve flow (*arrows*) and a second-degree atrioventricular block. The atrial contraction wave (A) is 140 bpm, and the ventricular contraction wave (V) is 70 bpm

The online version of the original article can be found under doi:[10.1007/s00246-008-9294-0](https://doi.org/10.1007/s00246-008-9294-0).

V. Tomek (✉) · J. Skovranek · R. A. Gebauer
Kardiocentrum and Cardiovascular Research Centre,
University Hospital Motol, V uvalu 84, Prague 5 150 08,
Czech Republic
e-mail: viktor.tomek@seznam.cz