Reply

Feico J. Halbertsma • Peter Andriessen

Received: 6 March 2009 / Accepted: 22 March 2009 / Published online: 15 April 2009
© Springer-Verlag 2009

Sir,

In the case presented by us showing cerebral air after cardiopulmonary resuscitation (CPR), we speculated on the cause of sudden cardiac arrest [1]. An occluding thrombus in the inferior vena cava found on post-mortem examination was identified as a likely initial cause, and we considered the vascular air entry as secondary to CPR. In a large series reported by Shiotani et al. [2] of 228 resuscitated patients without a history of pulmonary interstitial emphysema (PIE), 71% showed some degree of intravascular air.

Royal and Merrow [3] in their letter suggest that lymphatic drainage of PIE may have caused the cardiac arrest and may explain the venous vascular air. Indeed, the neonate we reported had been ventilated due to idiopathic respiratory distress syndrome, which had been complicated by temporary pneumothorax. However, the stable clinical condition over 5 days with spontaneous ventilation without any respiratory symptoms or need for supplemental oxygen makes the presence of PIE unlikely. Thus PIE as the initial cause of cardiac arrest is improbable, whereas PIE secondary to CPR is, in our opinion, very well possible due to both high inflation and chest compression pressures. Both the traditional assumed mechanism (traumatic alveolar-vascular fistula), and the mechanism proposed by Royal and Merrow (vascular air entry of PIE through lymphatic vessels) based on earlier studies by Booth et al. [4] offer likely pathways for vascular air entry, and also for

predominant venous air localization—the former mechanism through backward flow by CPR, the latter through antegrade flow after entry into the venous system through the right lymphatic and venous lymphatic duct. Indeed, we noted venous air content exceeding arterial air content, and right ventricle air exceeding left ventricle air based on calculations using data from post-mortem whole-body HRCT (total intravascular air estimated at 9.5 ml).

We still consider that the clinical course was more consistent with vascular air entry secondary to CPR, similar to the observations of Shiotani et al. [2]; however, preexisting PIE cannot be ruled out and is another possible cause.

References

1. Halbertsma FJ, Andriessen P (2009) Cerebral air embolism in a neonate following cardiopulmonary resuscitation. *Pediatr Radiol* 39:92
2. Shiotani S, Kohno M, Ohashi N et al (2005) Cardiovascular gas on non-traumatic postmortem computed tomography (PMCT): the influence of cardiopulmonary resuscitation. *Radiat Med* 23: 225–229
3. Royal SA, Merrow AC Jr (2009) Cerebral air embolism in a neonate following cardiopulmonary resuscitation. *Pediatr Radiol*. doi:10.1007/s00247-009-1196-7
4. Booth TN, Royal SA, Allen BA (1995) Lymphatic air embolism: a new hypothesis regarding the pathogenesis of neonatal systemic air embolism. *Pediatr Radiol* 25:S220–S227

F. J. Halbertsma (✉) • P. Andriessen
Neonatal Intensive Care Unit,
Department of Pediatrics and Neonatology,
Máxima Medical Center,
P.O. Box 7777, 5500 MB Veldhoven, The Netherlands
e-mail: F.Halbertsma@mmc.nl