Whole-body MR and cysticercosis

Viroj Wiwanitkit

Received: 27 January 2010 / Accepted: 12 February 2010
© Springer-Verlag 2010

Sir,

I read the report on cysticercosis by Kumar et al. [1] with great interest. Kumar et al. concluded that “Whole-body MR diagnosis of an infectious disease is unique” [1]. Indeed, whole-body MR diagnosis is useful for cysticercosis management, but it is not totally specific to this disease. There are also other cystic parasitic diseases in which whole-body MR diagnosis is useful and the finding of disseminated cysts can also be seen. The disseminated hydatid cyst is the best example [2, 3]. This cystic disease should be an important differential diagnosis for cysticercosis.

References

1. Kumar A, Goenka AH, Choudhary A et al (2010) Disseminated cysticercosis in a child: whole-body MR diagnosis with the use of parallel imaging. *Pediatr Radiol* 40:223–227
2. Moharamzad Y, Kharazi HH, Shobeiri E et al (2008) Disseminated intraspinal hydatid disease. *J Neurosurg Spine* 8:490–493
3. Dziri C, Haouet K, Fingerhut A et al (2009) Management of cystic echinococcosis complications and dissemination: where is the evidence? *World J Surg* 33:1266–1273

V. Wiwanitkit (✉)
Wiwanitkit House,
Bangkhuae,
10160 Bangkok, Thailand
e-mail: wviroj@yahoo.com