Erratum to: Pre-medication to block [^{18}F]FDG uptake in the brown adipose tissue of pediatric and adolescent patients

Michael J. Gelfand · Sara M. O'Hara ·
Lois A. Curtwright · Joseph R. MacLean

Published online: 1 February 2011
© Springer-Verlag 2011

Erratum to: Pediatr Radiol
DOI 10.1007/s00247-005-1505-8

Table 1 contains erroneous data. A corrected table is presented here.

Table 1 Pre-medication used to block brown adipose tissue uptake of [^{18}F]FDG

<i>N</i>	Drug and route	Dosage	Mean dose	Maximum dose
34	Diazepam, p.o. (low dose)	<0.08 mg/kg	0.058 mg/kg	
9	Diazepam, p.o. (moderate dose)	>=0.08 mg/kg	0.096 mg/kg	7.5 mg
45	Fentanyl, i.v.	<= 1.0 mcg/kg	0.75 mcg/kg	50 mcg
23	No pre-medication			
7	On opiates for pain, by any route of administration			

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00247-005-1505-8>.

M. J. Gelfand (✉) · S. M. O'Hara · L. A. Curtwright ·
J. R. MacLean
Department of Radiology,
Cincinnati Children's Hospital Medical Center,
3333 Burnet Ave,
Cincinnati, OH 45229-3039, USA
e-mail: Michael.gelfand@cchmc.org