An informative set of SSLP markers and genomic profiles in the rat MHC, the *RT1* complex

Yumie Takagi · Takashi Kuramoto · Birger Voigt ·
Toshiko Tsurumi · Satoshi Nakanishi ·
Tomoji Mashimo · Norio Masui · Tadao Serikawa

Published online: 2 April 2009
© Springer-Verlag 2009

Erratum to: Immunogenetics
DOI 10.1007/s00251-008-0352-9

Unfortunately, Figure 1 was published with errors. The correct Figure 1 is given below.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00251-008-0352-9>.

Y. Takagi · T. Kuramoto · B. Voigt · T. Tsurumi · S. Nakanishi ·
T. Mashimo · T. Serikawa (✉)
Institute of Laboratory Animals, Graduate School of Medicine,
Kyoto University,
Yoshidakonoe-cho Sakyo-ku,
Kyoto 606-8501, Japan
e-mail: serikawa@anim.med.kyoto-u.ac.jp

Y. Takagi · N. Masui
Japan SLC, Inc.,
Shizuoka, Japan

their flanking genes (Hurt et al. 2004). SSLP marker profiles of 67 rat strains. These rat strains are sorted based on the assignment on the phylogenic tree. Same haplotype patterns are individually colored. The SSLP marker haplotype of the *RT1* complex is roughly divided into three genomic regions: 1; *D20Rat47* to *D20Yum29*, 2; *D20Yum84* to *D20Yum41*, 3; *D20Yum42* to *D20Yum82*