Erratum to: *Lactobacillus reuteri* CRL 1101 highly produces mannitol from sugarcane molasses as carbon source

Maria Eugenia Ortiz · María José Fornaguera ·
Raúl R. Raya · Fernanda Mozzi

Published online: 4 August 2012
© Springer-Verlag 2012

Erratum to: Appl Microbiol Biotechnol DOI 10.1007/s00253-012-3945-z

The published version of this article unfortunately contains mistakes. Tables 1 and 2 have error numbers; the standard

deviation (SD) values are higher than the real ones as a number 3 was added incorrectly at the beginning of each SD value in all cases, except for the SD corresponding to “Mol 3 %, 37 °C” in Table 1, which is fine. The correct Tables 1 and 2 are the following:

Table 1 Stoichiometric parameters for mannitol and organic acid formation by *L. reuteri* CRL 1101 using sugarcane molasses as substrate in MRS-supplemented media after 24 h fermentations

Culture conditions	Consumption (mMol)		Mtl ^c (mMol)	Y _{Mtl} %	Mtl/HLac ^d	Mtl/HLac + HAc ^c
	Glucose	Fructose				
Mol ^a 3 %, 37 °C	89.1±3.0	89.2±4.3	67.3±12.7	75.5	0.84	0.59
Mol 5 %, 37 °C (w/v) mol	169.0±22.2	168.0±21.2	108.5±25.4	64.6	0.77	0.56
Mol 7.5 %, 37 °C	174.1±35.8	198.0±38.2	177.6±26.6	89.7	1.03	0.75
Mol 10 %, 37 °C	174.6±19.2	208.0±23.2	184.5±22.5	88.6	1.08	0.73
Mol 7.5 %, 30 °C	67.9±2.8	59.0±12.7	48.0±9.7	82.3	0.98	0.64
Mol 7.5 %, 37 °C A ^b	201.4±10.0	243.0±5.2	211.3±15.5	86.9	1.24	0.83

Mtl mannitol, Mtl/HLac ratio of mannitol to lactic acid, Mtl/HLac + HAc mannitol ratio to lactic acid + acetic acid

^a Molasses

^b Agitated cultures

Maria Eugenia Ortiz and María José Fornaguera contributed equally to the article.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00253-012-3945-z>.

M. E. Ortiz · M. J. Fornaguera · R. R. Raya · F. Mozzi (✉)
Centro de Referencia para Lactobacilos (CERELA)-CONICET,
Chacabuco 145,
San Miguel de Tucumán 4000, Argentina
e-mail: fmozzi@cerela.org.ar

Table 2 Mannitol 2-dehydrogenase activity of cell-free extracts of *L. reuteri* CRL 1101 grown in molasses-supplemented MRS medium under different culture conditions

Culture conditions	Specific MDH activity (U/mg cell protein)			
	Time of incubation (h)			
	4	8	24	48
Mol ^a 3 %, 37 °C	3.646±0.052	2.100±0.097	2.340±0.024	2.016±0.067
Mol 5 %, 37 °C	2.587±0.098	1.997±0.257	2.122±0.134	1.960±0.086
Mol 7.5 %, 37 °C	2.277±0.036	1.847±0.144	1.726±0.308	1.580±0.081
Mol 10 %, 37 °C	2.598±0.079	2.094±0.662	2.087±0.083	1.349±0.089
Mol 7.5 %, 30 °C	0.260±0.086	2.712±0.646	2.685±0.385	1.743±0.075
Mol 7.5 %, 37 °C A ^b	2.064±0.145	1.037±0.320	1.075±0.291	0.583±0.166

^aMolasses

^bAgitated culture