



Correction to: Metabolic pathway of 6-aminohexanoate in the nylon oligomer-degrading bacterium *Arthrobacter* sp. KI72: identification of the enzymes responsible for the conversion of 6-aminohexanoate to adipate

Ikki Takehara¹ · Tsubasa Fujii¹ · Yuuki Tanimoto¹ · Dai-ichiro Kato² · Masahiro Takeo¹ · Seiji Negoro¹

Published online: 12 December 2017

© Springer-Verlag GmbH Germany, part of Springer Nature 2017

Correction to: Appl Microbiol Biotechnol

<https://doi.org/10.1007/s00253-017-8657-y>

The original publication of this paper contains mistakes. Below the needed corrections:

Table 1 legend:

The following words "In Tables 1 and 2," are added at the beginning of the last sentence of the legend of the table. "Enzyme" is changed to "enzyme".

Table 2 legend:

Last sentence "Enzyme samples obtained by independent purification experiments were used" is deleted.

Figures:

The sublabels "A", "B" to small letters "a", "b", for Figs. 2, 4, 5, 6, and 7.

The original article has been corrected.

The online version of the original article can be found at <https://doi.org/10.1007/s00253-017-8657-y>.

✉ Seiji Negoro
negoro@eng.u-hyogo.ac.jp

¹ Department of Applied Chemistry, Graduate School of Engineering, University of Hyogo, 2167 Shosha, Himeji, Hyogo 671-2280, Japan

² Department of Chemistry and Bioscience, Graduate School of Science and Engineering, 1-21-35 Korimoto, Kagoshima, Kagoshima 890-0065, Japan