N. Subba Rao
I. Saroja Nirmala
K. Suryanarayana

Groundwater quality in a coastal area: a case study from Andhra Pradesh, India

Published online: 4 January 2006
© Springer-Verlag 2006

The online version of the original article
can be found at <http://dx.doi.org/10.1007/s00254-005-1306-2>

Environ Geol (2005). DOI: 10.1007/s00254-005-1306-2

In the online version in the second paragraph below the heading “Factors controlling groundwater quality” and in the print version on page 548, 2nd column, 2nd paragraph, the first sentence should read:

Though the groundwater samples 1, 2, 3, 12 and 19 show both fresh,

TDS < 1,000 mg/l, (sample 12) and brackish, TDS > 1,000 mg/l, (samples 1–3 and 19) categories (Table 1), the ratios of $\text{Ca}^{2+}:\text{Mg}^{2+} < 1$ (sample 19), $\text{TA}:\text{TH} < 1$ (samples 1–3 and 12) and $\text{Cl}:\text{HCO}_3^- > 1$ (samples 1–3 and 12) in groundwaters (Table 3) suggest the influence of seawater.

N. Subba Rao (✉) · I. Saroja Nirmala
K. Suryanarayana
Hydrogeology Laboratory,
Department of Geology,
Andhra University,
Visakhapatnam 530 003, India
E-mail: srnandipati@rediffmail.com
Tel.: +91-0891-2844716
Fax: +91-0891-2755324