Sylvain R. Duc
Klaus U. Wentz
Kurt P. Käch
Christoph L. Zollikofer

First report of an accessory popliteal muscle: detection with MRI

Published online: 10 August 2004
© ISS 2004

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00256-004-0775-9>

S. R. Duc · K. U. Wentz (✉) · C. L. Zollikofer
MR Research Group of the Institute of Diagnostic Radiology,
Kantonsspital Winterthur,
Brauerstrasse 15, 8401 Winterthur, Switzerland
e-mail: Ku.wentz@ksw.ch
Tel.: +41-52-2664610
Fax: +41-52-2664603

K. P. Käch
Department of Trauma Surgery,
Kantonsspital Winterthur, Winterthur, Switzerland

Skeletal Radiol (2004) 33:429–431

The following abbreviations, which were found in Figures 1 and 3 were previously undefined:

BF Biceps femoris, *Fe* Femur, *FI* Fibula, *G* Gracilis, *GM* Gastrocnemius medialis, *GL* gastrocnemius lateralis, *LFC* Lateral femoral condyle, *MFC* Medial femoral condyle, *MTP* Medial tibial plateau, *P* Popliteal artery, *PA* Popliteus accessorius, *Pop* Popliteus, *S* Sartorius, *SM* Semimembranosus, *ST* Semitendinosus, *TI* Tibia

Fig. 1 Axial image shows the location of the accessory popliteus muscle deep in the popliteal fossa, passing between the posterior articular capsule and the popliteal artery

Fig. 3 Coronal image nicely shows the oblique descending course of the accessory popliteal muscle from the lateral femoral condyle towards the medial femorotibial compartment