ERRATUM

Erratum to: Imaging findings of anterior hip dislocations

Kyle Pfeifer¹ · Michael Leslie² · Kirsten Menn³ · Andrew Haims³

Published online: 13 March 2017
© ISS 2017

Erratum to: Skeletal Radiol
DOI 10.1007/s00256-017-2605-x

Figure captions 9 and 10 were reversed in the original article.
The correct figures and captions are indicated below:
The original article was corrected.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00256-017-2605-x>.

✉ Kyle Pfeifer
kpfeife@gmail.com

Michael Leslie
Michael.leslie@yale.edu

Kirsten Menn
kirsten.menn@gmail.com

Andrew Haims
Andrew.haims@yale.edu

¹ Department of Radiology, Mallinckrodt Institute of Radiology, 510 South Kings Highway Boulevard, St. Louis, MO, USA

² Department of Orthopedics and Rehabilitation, Yale School of Medicine, 47 College Place, New Haven, CT 06520, USA

³ Department of Radiology and Biomedical Imaging, Yale University School of Medicine, 333 Cedar St, New Haven, CT 06520, USA

Fig. 9 A 27-year-old female status post motor vehicle accident. Figures are surface-rendered 3D images from CT of the pelvis using (Vitrea®, Minnetonka, MN). **a** An anterior inferior dislocation of the

femoral head (*red arrows*) with the head resting over the left obturator foramen. **b** The patient's femoral head lodged within the obturator foramen

Fig. 10 Radiographs are 0 months (A: Z = 0.48), 5 months (B: Z = 0.48), following an anterior inferior right hip dislocation. **a** Post reduction image. Five months later, the patient returned with pain. **b** Osteonecrosis and subchondral collapse (*red arrows*). Unfortunately,

the patient had pain with ambulation and subsequently underwent right total hip arthroplasty. Incidentally noted is a benign-appearing mixed sclerotic and lucent lesion in the proximal femur