Beta-irradiation used for systemic radioimmunotherapy induces apoptosis and activates apoptosis pathways in leukaemia cells

Claudia Friesen¹, Annelie Lubatschowski¹, Jörg Kotzerke², Inga Buchmann², Sven N. Reske², Klaus-Michael Debatin¹

¹ University Children's Hospital, Ulm, Germany

² Department of Nuclear Medicine, University of Ulm, Ulm, Germany

Published online: 26 July 2003

© Springer-Verlag 2003

Eur J Nucl Med Mol Imaging (2003) 30:1262

DOI 10.1007/s00259-003-1314-y

The online version of the original article can be found at
<http://dx.doi.org/10.1007/s00259-003-1216-z>

Eur J Nucl Med Mol Imaging (2003)

DOI 10.1007/s00259-003-1216-z

In the Materials and methods section, under the heading “Western blot analysis”, “rabbit anti-Bcl-X_{3/2}” should read “rabbit anti-Bcl-x_{S/L}”.

Klaus-Michael Debatin (✉)

University Children's Hospital, Prittwitzstrasse 43, 89075 Ulm, Germany

e-mail: klaus-michael.debatin@medizin.uni-ulm.de

Tel.: +49-731-50027700, Fax: +49-731-50026681