A new approach for improving diagnostic accuracy in Alzheimer's disease and frontal lobe dementia utilising the intrinsic properties of the SPET dataset

Marco Pagani^{1, 2}, Vassili A. Kovalev^{3, 4}, Roger Lundqvist⁵, Hans Jacobsson^{2, 6}, Stig A. Larsson², Lennart Thurfjell⁵

¹ Institute of Cognitive Sciences and Technology, CNR, Rome, Italy

² Section of Nuclear Medicine, Karolinska Hospital/Institute, Stockholm, Sweden

³ Institute of Engineering Cybernetics, Belarus National Academy of Sciences, Minsk, Belarus

⁴ Max-Planck Institute of Cognitive Neuroscience, Leipzig, Germany

⁵ Applied Medical Imaging, Uppsala, Sweden

⁶ Department of Radiology, Karolinska Hospital, Stockholm, Sweden

Published online: 19 December 2003

© Springer-Verlag 2003

Eur J Nucl Med Mol Imaging (2004) 31:307

DOI 10.1007/s00259-003-1434-4

Marco Pagani (✉)

Institute of Cognitive Sciences and Technology, CNR, Rome, Italy

e-mail: marco_00198@yahoo.com

Fax: +39-06-824737

Eur J Nucl Med Mol Imaging (2003)

30:1481–1488

The first affiliation of Marco Pagani was omitted by mistake in the printed version. Please see the complete and correct affiliations above.

The online version of the original article can be found at

<http://dx.doi.org/10.1007/s00259-003-1196-z>