Myocardial metabolism of ^{123}I -BMIPP under low-dose dobutamine infusion: implications for clinical SPECT imaging of ischemic heart disease

Ryohei Hosokawa¹, Ryuji Nohara², Taku Hirai⁴, Yasuhisa Fujibayashi³, Masatoshi Fujita¹, Naoshige Kambara¹, Muneo Ohba¹, Eiji Tadamura¹, Takeshi Kimura¹, Toru Kita¹

¹ Department of Cardiovascular Medicine, and Nuclear Medicine, Kyoto University Graduate School of Medicine, Kyoto, Japan

² Department of Cardiology, Kitano Hospital, The Tazuke Kofukai Medical Research Institute, 2-4-20 Ohgimachi, Kita-ku, Osaka 530-8480, Japan

³ Molecular Imaging, Biomedical Imaging Research Center, Fukui Medical University, Fukui, Japan

⁴ Department of Cardiology, Nara Hospital, Kinki University School of Medicine, Nara, Japan

Published online: 18 March 2005

© Springer-Verlag 2005

Eur J Nucl Med Mol Imaging (2005) 32:520
DOI 10.1007/s00259-005-1782-3

The online version of the original article can be found at
<http://dx.doi.org/10.1007/s00259-004-1622-x>

Ryuji Nohara (✉)
Department of Cardiology, Kitano Hospital
2-4-20 Ohgimachi, Kita-ku, Osaka 530-8480, Japan
e-mail: noharar@kitano-hp.or.jp
Tel.: +81-6-63128824, Fax: +81-6-63128867

Eur J Nucl Med (2005) 32:75–83

A misprint occurred in the right-hand column of page 77, partly obscuring the equation. The equation is reproduced here.

$$\text{Extraction fraction} = 1 - \frac{\left[\begin{smallmatrix} ^{123}\text{I in the blood (0 to 30s)} \end{smallmatrix} \right] / \left[\begin{smallmatrix} ^{123}\text{I injected} \end{smallmatrix} \right]}{\left[\begin{smallmatrix} ^{125}\text{I in the blood (0 to 30s)} \end{smallmatrix} \right] / \left[\begin{smallmatrix} ^{125}\text{I injected} \end{smallmatrix} \right]}$$