Prognostic study of risk stratification among Japanese patients with ischemic heart disease using gated myocardial perfusion SPECT: J-ACCESS study

Tsunehiko Nishimura • Kenichi Nakajima •
Hideo Kusuoka • Akira Yamashina •
Shigeyuki Nishimura

Published online: 26 July 2008
© Springer-Verlag 2008

Erratum to: Eur J Nucl Med Mol Imaging
DOI 10.1007/s00259-007-0608-x

Eur J Nucl Med (2008) 35:319–328

In Table 1, the entry for “Family history of coronary artery disease” should read “394/3,325 (11.9%)” instead of “1,871/3,968 (47.2%)”.

The same data were mistakenly copied from the line above.

The online version of the original can be found at <http://dx.doi.org/10.1007/s00259-007-0608-x>.

T. Nishimura (✉)
Department of Radiology, Graduate School of Medical Science,
Kyoto Prefectural University of Medicine,
465 Kajicho, Kawara-machi, Hirokoji,
Kamigyo-ku 602-8566 Kyoto, Japan
e-mail: nishimura@koto.kpu-m.ac.jp

H. Kusuoka
Osaka National Hospital,
Osaka, Japan

A. Yamashina
Second Department of Internal Medicine,
Tokyo Medical University Hospital,
Tokyo, Japan

S. Nishimura
Division of Cardiology, Saitama Medical School Hospital,
Saitama, Japan

K. Nakajima
Department of Biotracer Medicine, Kanazawa University
Graduate School of Medical Science,
Kanazawa, Japan
e-mail: nakajima@med.kanazawa-u.ac.jp