Erratum to: Impaired myocardial blood flow reserve in subjects with metabolic syndrome analyzed using positron emission tomography and N-13 labeled ammonia

Hiroki Teragawa · Koichi Morita · Hiroki Shishido ·
Nobuaki Otsuka · Yutaka Hirokawa ·
Kazuaki Chayama · Nagara Tamaki · Yasuki Kihara

Published online: 15 July 2010
© Springer-Verlag 2010

Erratum to: Eur J Nucl Med Mol Imaging
DOI 10.1007/s00259-009-1307-6

Table 3 contained some incorrect data. The correct table is reproduced here in full.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s00259-009-1307-6>.

H. Teragawa (✉) · Y. Kihara
Department of Cardiovascular Medicine,
Hiroshima University Graduate School of Biomedical Sciences,
1-2-3 Kasumi,
Minami-ku, Hiroshima 734-8551, Japan
e-mail: hteraga@gmail.com

K. Morita · N. Tamaki
Department of Nuclear Medicine,
Hokkaido University Graduate School of Medicine,
Sapporo, Japan

H. Shishido · N. Otsuka · Y. Hirokawa
Hiroshima Heiwa Clinic,
Hiroshima, Japan

K. Chayama
Department of Molecular Science and Medicine,
Hiroshima University Graduate School of Biomedical Sciences,
Hiroshima, Japan

Table 3 Relationship between MFR and other parameters

Univariate analyses			Multivariate analyses		
Factor	MFR		Factor	MFR	
	r	p Value		F value	p Value
Abdominal obesity	.	0.1828	HOMA-IR	4.7226	0.0408
Elevated triglycerides	.	0.3596	Hypertension	4.5855	0.0436
Low HDL-cholesterolemia	.	0.8144	Posterior wall thickness	2.9044	0.1024
Hypertension	.	0.0255	Body mass index	0.5697	0.4584
Hyperglycemia	.	0.1377			
Taking statin	.	0.0921			
Body mass index	-0.444	0.0204			
Waist circumference	-0.353	0.0705			
Fasting blood sugar	-0.359	0.0655			
HOMA-IR	-0.507	0.0075			
Hemoglobin A1C	-0.363	0.0624			
Interventricular septal thickness	-0.367	0.0595			
Posterior wall thickness	-0.417	0.0303			
Visceral fat area	-0.361	0.0651			

HOMA-IR, homeostasis model assessment-insulin resistance; LDL, low-density lipoprotein; MFR, myocardial flow reserve. $r^2=0.449$