Kinetics of 3-[¹²³I]iodo-L-α-methyltyrosine transport in rat C6 glioma cells

Burkhard Riemann¹, Florian Stögbauer², Klaus Kopka¹, Hartmut Halfter², Manuela Lasic¹, Anja Schirmacher², Torsten Kuwert³, Matthias Weckesser¹, E. Bernd Ringelstein², Otmar Schober¹

¹ Department of Nuclear Medicine, Westfälische Wilhelms-Universität Münster, Albert-Schweitzer-Strasse 33, 48129 Münster, Germany

² Department of Neurology, Westfälische Wilhelms-Universität Münster, Germany

³ Department of Nuclear Medicine, Friedrich-Alexander-Universität Erlangen, Germany

Published online: 10 November 2000

© Springer-Verlag 2000

Eur J Nucl Med (2000) 27:1870

DOI 10.1007/s002590000422

Eur J Nucl Med (1999) 26:1274–1278

The fourth complete paragraph of column 2 on p. 1275 stated that “After addition of a surplus of unlabelled IMT to the medium, thereby saturating all binding sites of the specific transport system for ¹²³I-IMT, less than 0.1% unspecific uptake of radiolabelled ¹²³I-IMT into the cells was found.” In fact, “less than 0.1%” should have read “less than 10%”.