ERRATUM

Erratum to: Perfusion changes in gastric adenocarcinoma: evaluation with 64-section MDCT

Jin Yao,¹ Zhi-gang Yang,^{1,2} Tian-wu Chen,¹ Yuan Li,¹ Lin Yang¹¹Department of Radiology, West China Hospital, Sichuan University, 37# Guo Xue Xiang, Chengdu, Sichuan 610041, China²National Key Laboratory of Biotherapy Center, West China Hospital, Sichuan University, 37# Guo Xue Xiang, Chengdu, Sichuan 610041, China

Erratum to: Abdom Imaging (2010) 35:195–202 DOI 10.1007/s00261-009-9503-1

In the Tables 1, 2, 3, and 5, the unit of perfusion was stated incorrectly as mL/100 g/min. It should have been stated as mL/min/mL. The corrected tables are as follows:

Table 1. Intraobserver variability for PF, PE, TTP, and BV measurement of normal stomach

Parameter measurement	ICC ^a (95% CI)	Mean difference	SD	95% limits of agreement
PF (mL/min/mL)	0.929 (0.894–0.952)	−1.44	13.69	−28.27, 25.38
PE (Hu)	0.969 (0.954–0.979)	−0.29	8.24	−16.45, 15.87
TTP (s)	0.816 (0.726–0.877)	0.21	10.22	−19.83, 20.25
BV (mL/100 g)	0.936 (0.905–0.957)	0.24	5.81	−11.14, 11.62

^aICC, intraclass correlation coefficient**Table 2.** Intraobserver variability for PF, PE, TTP, and BV measurement of gastric adenocarcinoma

Parameter measurement	ICC ^a (95% CI)	Mean difference	SD	95% limits of agreement
PF (mL/min/mL)	0.922 (0.868–0.954)	2.38	8.05	−13.39, 18.15
PE (Hu)	0.990 (0.984–0.994)	0.10	2.25	−4.30, 4.51
TTP (s)	0.969 (0.948–0.982)	−1.21	4.20	−9.45, 7.03
BV (mL/100 g)	0.968 (0.947–0.981)	−0.68	5.23	−10.94, 9.58

^aICC, intraclass correlation coefficient

The online version of the original article can be found under doi:[10.1007/s00261-009-9503-1](https://doi.org/10.1007/s00261-009-9503-1).

Correspondence to: Zhi-gang Yang; email: yangzg1117@yahoo.com.cn

Table 3. Perfusion values in gastric adenocarcinoma vs. in normal stomach

Location	Perfusion parameter	Tumor ^a	Normal stomach ^a	<i>P</i> value
Cardia	PF (mL/min/mL)	28.87 ± 11.50	24.58 ± 20.91	0.45
	PE (Hu)	24.51 ± 8.94	20.55 ± 11.49	0.20
	TTP (s)	29.98 ± 13.12	31.42 ± 15.22	0.55
	BV (mL/100 g)	19.78 ± 21.02	12.35 ± 12.93	0.14
Body	PF (mL/min/mL)	31.73 ± 19.56	40.98 ± 30.29	0.19
	PE (Hu)	29.49 ± 11.70	35.28 ± 29.95	0.48
	TTP (s)	31.28 ± 11.54	30.18 ± 8.89	0.75
	BV (mL/100 g)	16.02 ± 7.02	15.47 ± 11.13	0.83
Antrum and pylorus	PF (mL/min/mL)	28.02 ± 14.96	30.35 ± 14.40	0.59
	PE (Hu)	28.67 ± 15.87	29.02 ± 12.23	0.93
	TTP (s)	34.77 ± 12.23	34.29 ± 11.71	0.89
	BV (mL/100 g)	20.80 ± 12.41	11.29 ± 9.99	0.006
Whole stomach	PF (mL/min/mL)	30.28 ± 14.89	32.85 ± 25.69	0.49
	PE (Hu)	27.89 ± 11.29	29.06 ± 23.30	0.72
	TTP (s)	32.60 ± 12.30	31.46 ± 11.86	0.57
	BV (mL/100 g)	19.75 ± 14.74	13.59 ± 11.46	0.004

^aData are means ± standard deviations**Table 5.** Perfusion values of subgroups with lymph node metastases vs. without lymph node metastases

Perfusion parameter	With lymph node metastases (<i>n</i> = 29) ^a	Without lymph node metastases (<i>n</i> = 13) ^a	<i>P</i> value
PF (mL/min/mL)	30.67 ± 12.36	31.10 ± 21.35	0.93
PE (Hu)	29.32 ± 9.90	25.44 ± 11.17	0.29
TTP (s)	32.58 ± 11.99	32.55 ± 16.51	0.99
BV (mL/100 g)	18.75 ± 9.89	15.62 ± 11.15	0.39

^aData are means ± standard deviations