The “watering can perineum”

Colin M. Segovis , Raymond B. Dyer

Department of Radiology, Wake Forest Baptist Medical Center, Wake Forest University School of Medicine, Medical Center Blvd, Winston-Salem, NC 27157, USA

A watering can (Fig. 1) metaphorically describes the fluoroscopic demonstration of contrast opacifying multiple fistulas from the urethra to the perineum, as seen on voiding or retrograde urethrography (Fig. 2) [1]. In patients with this condition, urine exits through multiple perineal openings, hence the descriptive “watering can” [2]. The urethroperineal fistulas are sequelae of traumatic or chronic inflammatory urethral strictures [2]. Classically, this sign is associated with advanced infection of the urethra and adjacent tissues, most commonly gonorrhea. Associated paraurethral abscesses are frequent [1]. Other etiologies include schistosomiasis, tuberculosis, and Crohn’s disease [1, 2].

The radiologic appearance is specific to fistula formation rather than a particular etiology. Consequently, determination of the cause of the fistula formation requires an appropriate history and urinalysis, and possibly urethroscopy and biopsy [1].

Fig. 1. The watering can disperses the water stream through multiple openings in the head.

Fig. 2. A retrograde urethrogram opacifies multiple urethroperineal fistulas in a patient with a “watering can perineum”.

Compliance with ethical standards

Ethical approval This article does not contain any studies with animals or human participants performed by any of the authors.

Conflict of Interest Dr. Segovis declares that he has no conflict of interest. Dr. Dyer declares that he has no conflict of interest.

References

1. Friedland G, Nino-Murcia M, Gill H, Perkas I (1998) Acquired nontraumatic lesions of the urethra. In: Jafri S, Diokno A, Amendola M (eds) *Lower genitourinary radiology: imaging and intervention*, 1st edn. New York: Springer, pp 352–389
2. Zagoria RJ, Dyer RB, Brady C (2016) *Genitourinary imaging: the requisites*. Philadelphia: Elsevier Inc.