Erratum to: Hepatocellular carcinoma detection: diagnostic performance of a simulated abbreviated MRI protocol combining diffusion-weighted and T1-weighted imaging at the delayed phase post gadoxetic acid

Cecilia Besa,^{1,2} Sara Lewis,² Pari V. Pandharipande,³ Jagpreet Chhatwal,³ Amita Kamath,² Nancy Cooper,² Ashley Knight-Greenfield,¹ James S. Babb,⁴ Paolo Boffetta,⁵ Norma Padron,⁶ Claude B. Sirlin,⁷ Bachir Taouli^{1,2}

¹Translational and Molecular Imaging Institute, Icahn School of Medicine at Mount Sinai, 1470 Madison Avenue, New York, NY 10029, USA

²Department of Radiology, Icahn School of Medicine at Mount Sinai, New York, NY 10029, USA

³Institute for Technology Assessment, Department of Radiology, Massachusetts General Hospital, Boston, MA, USA

⁴Department of Radiology, New York University Langone Medical Center, New York, NY, USA

⁵Division of Cancer Prevention and Control, Department of Oncological Sciences, Icahn School of Medicine at Mount Sinai, New York, NY, USA

⁶Department of Population Health Science and Policy, Icahn School of Medicine at Mount Sinai, New York, NY, USA

⁷Liver Imaging Group, Department of Radiology, University of California, San Diego, CA, USA

Erratum to: Abdom Radiol (2017) 42: 179–190 DOI 10.1007/s00261-016-0841-5

In the original version of the article, the reference number 11 was incorrectly published. It has been corrected now. The correct reference number 11 is given below:

Reference

11. Marks RM, Ryan A, Heba ER, et al. (2015) Diagnostic per-patient accuracy of an abbreviated hepatobiliary phase gadoxetic acid-enhanced MRI for hepatocellular carcinoma surveillance. *AJR Am J Roentgenol* 204(3):527–535

The online version of the original article can be found under doi:
[10.1007/s00261-016-0841-5](https://doi.org/10.1007/s00261-016-0841-5).

Correspondence to: Bachir Taouli; email: bachir.taouli@mountsinai.org