Correction to: Differentiating focal nodular hyperplasia from hepatocellular adenoma: Is hepatobiliary phase MRI (HBP-MRI) using linear gadolinium chelates always useful?

Marion Roux¹,^{1b} Frederic Pigneur,² Laurence Baranes,^{2,3} Julien Calderaro,^{4,3} Mélanie Chiaradia,^{2,3} Thomas Decaens,^{5,3} Sandrine Kastahian,⁶ Anaïs Charles-Nelson,⁶ Lambros Tselikas,² Charlotte Costentin,^{5,3} Alexis Laurent,^{7,3} Daniel Azoulay,^{7,3} Ariane Mallat,^{5,3} Alain Rahmouni,^{2,3} Alain Luciani^{2,3,8}

¹Service de Radiodiagnostic et Radiologie interventionnelle, CHUV, Rue du Bugnon 46, 1011 Lausanne, Switzerland

²Groupe Henri Mondor Albert Chenevier, Imagerie Medicale, AP-HP, 94010 Creteil, France

³Faculte de Medecine de Creteil, Universite Paris Est Creteil, 94010 Creteil, France

⁴Groupe Henri Mondor Albert Chenevier, Pathology, AP-HP, 94010 Creteil, France

⁵Groupe Henri Mondor Albert Chenevier, Hepato-Gastroenterology Department, AP-HP, 94010 Creteil, France

⁶Groupe Henri Mondor Albert Chenevier, Unite de Recherche Clinique (URC), AP-HP, 94010 Creteil, France

⁷Groupe Henri Mondor Albert Chenevier, Liver Surgery, AP-HP, 94010 Creteil, France

⁸INSERM Unite U 955, Equipe 18, 94010 Creteil, France

Correction to: Abdom Radiol (2017)

<https://doi.org/10.1007/s00261-017-1377-z>

The original version of this article unfortunately contained a mistake. There was an error in the last sentence of the summary and the last sentence of the discussion. It should read as “On HBP, all FNH appeared hyper- or iso-intense compared to adjacent liver while close to 97% of HCA appeared hypointense”.