ERRATUM

Christian M. Krawczyk · Srdan Verstovšek
Peter Ujházy · Darbie Maccubbin · M. Jane Ehrke

Protective specific immunity induced by cyclophosphamide plus tumor necrosis factor α combination treatment of EL4-lymphoma-bearing C57BL/6 mice

In: Cancer Immunol Immunother 40 (1995): 347–357

In the legends of both Fig. 5 (p 353) and Fig. 7 (p 354) of the above article the dosage for **CY (150 mg/kg)** have been omitted. The correct legends should read as follows:

Fig. 5 Thymic effector lytic activity – the effect of tumor rechallenge. Conditions were the same as those given in Fig. 2, **except the dose of Cy used was 150 mg/kg**. Mice were sacrificed on day 128 and thymocyte suspensions prepared. *R* mice rechallenged on day 60 by i.p. implantation of 5×10^4 viable EL4 tumor cells; *VR* mice not rechallenged. Evaluation of cytolytic activity after stimulation culture was against both EL4 or C1498 radiolabeled tumor target cells and in both 6-h and 18-h ^{51}Cr -release assays. *R* thymocytes were also examined for cytolytic activity on day 128. Immediately after the thymus had been obtained from the mice and single-cell suspensions prepared. These fresh *R* thymocytes (*F-R*) were plated with IL-2 (2.4 ng/ml) and radiolabeled target cells and ^{51}Cr release was assessed 18 h later

Fig. 7 Effect of combination treatment compared to CY alone on the thymic effector lytic activity. Conditions are the same as those given in Fig. 5, **where the dose of CY was 150 mg/kg**. Data using cells obtained on day 128 from rechallenged mice are shown