Clinical profile of long-term survivors of breast or thyroid cancer with metastatic spinal tumours

Y. Kasai · E. Kawakita · A. Uchida

Published online: 27 October 2006
© Springer-Verlag 2006

The online version of the original article can be found at: <http://dx.doi.org/10.1007/s00264-006-0145-4>.

Y. Kasai · E. Kawakita · A. Uchida
Department of Orthopaedic Surgery,
Mie University Graduate School of Medicine,
Tsu, Mie, Japan

Y. Kasai (✉)
2-174 Edobashi,
Tsu, Mie 514-8507, Japan
e-mail: ykasai@clin.medic.mie-u.ac.jp

Unfortunately, in Table 3, case 5, the condition of paraparesis is missing in the original article. The complete table is given here.

Table 3 Results of Tokuhashi's score

Case	Total score	Systemic condition	Number of spinal metastatic foci	Number of bone metastatic foci	Metastases in major organs	Condition of paraparesis (Frankel's classification)
1	15	Good	1	0	None	E
2	12	Mildly deteriorated	1	1	None	C
3	11	Severely deteriorated	2	0	None	C
4	10	Severely deteriorated	2	0	None	B
5	11	Severely deteriorated	1	0	None	B
6	13	Mildly deteriorated	1	1	None	D
7	8	Severely deteriorated	2	0	Lung	B
8	13	Good	2	1	None	D
9	15	Good	1	0	None	E
10	9	Severely deteriorated	1	0	Lung	B
11	10	Mildly deteriorated	1	1	None	C
12	8	Severely deteriorated	3	2	None	A
13	12	Mildly deteriorated	2	0	None	C
14	13	Good	2	1	None	E
15	9	Severely deteriorated	2	1	None	B
16	13	Mildly deteriorated	1	0	None	D
17	14	Good	1	0	None	D
18	8	Severely deteriorated	2	0	Brain	B
19	7	Severely deteriorated	4	0	Lung	A
20	11	Severely deteriorated	1	0	None	B
21	12	Mildly deteriorated	1	2	None	D
22	7	Severely deteriorated	2	1	Lung	B
23	12	Good	2	1	None	D
24	11	Severely deteriorated	1	0	None	B
25	14	Good	1	1	None	E