Letters to the Editor

A Safe and Effective Technique

I have recently read and digested the article **“External Ultrasound Used in Conjunction with Superficial Subdermal Liposuction: A Safe and Effective Technique,”** by authors Gasperoni, Salgarello and Gasperoni (*Aesthetic Plastic Surgery* 24:253–258, 2000). There are several things in the article which needs further explanation by the authors, mostly because many statements and conclusions are unsubstantiated in the article and leave the reader confused regarding the scientific grounds of the article. In other cases statements contradict each other and leave even more questions in readers’ minds. Let me take a few examples of things I would like to have explained further out of the article.

The article recommends using glycolic acid [30%] on our patients two to four weeks before surgery, applied topically on areas to be treated with liposuction. It is stated that this treatment will, “... smoothen the skin, remodel elastic fibers, and stimulate dermal collagen production, thus enhancing the skin retraction capability.” To my knowledge it seems incredible that this could be accomplished in a time of two to four weeks and, as I have been involved in a skin care clinic in my own practice for me last ten years. I would like to know more about the scientific bases for this statement.

We are also told that “when the skin is flabby a high protein diet or protein supplement may be suggested to shift the metabolic balance towards proteins in order to improve skin retraction.” Does this mean that a patient with a flabby skin, consulting for a facelift, could be treated with a high protein diet instead of surgery and therefore can expect a spontaneous skin retraction? As a plastic surgeon, I am, of course, interested in the scientific basis for the statement above.

We are recommended to treat our patients with vitamin C, 1000 mg per day for three months, as an antioxidant. However, in the discussion the authors state that external ultrasound “carries virtually no risk of free radical formation.” If there is no risk of free radical formation, why would one then prescribe an antioxidant for three months?

The authors also recommend keeping general anesthesia thoroughly deep in level, saying that other-

wise increased heart rate and blood pressure may cause bleeding, “This coagulation may thus be excessive, leading to uncontrolled bleeding and eventually to DIO.” This statement leads me to two thoughts: first, I have never heard of uncontrolled bleeding when doing a minimal liposuction like that the authors discuss and especially not when using a super-wet technique with lidocaine 0.05% and adrenaline. Second, most surgeons doing liposuction with local anesthesia alone would use this concentration of lidocaine for full analgesia of the area [unless internal ultrasound is used]. It eludes me why we would need deep general anesthesia in a case like this.

For local anesthesia, the authors are advocating a super-wet solution, which they define as when the volume of the wetting solution equals the estimated aspirate volume. They are using a concentration of lidocaine of 0.3% and recommending a maximum dosage of 20 mg per kg body weight which means that an ordinary woman of 60 kg could be infiltrated with a maximum of 400 ml of local anesthesia solution and subsequently, we would have an aspirate of maximum 400 ml, which makes me wonder what kind of case could be done under these circumstances.

When it comes to the act of delivering ultrasonic energy we are told to cover the probe filled with 2–3 cm of ultrasonic gel. It eludes me how any conduction of ultrasound can take place through such a thick layer of gel and I don’t really understand why it should be so thick. We are also told that when delivering the ultrasound, the skin becomes slightly warmer. However, further on in the article we are told that “external ultrasound does not significantly raise the skin temperature.” As a reader, I would like to know whether the skin temperature is raised by the treatment or not, and if so why?

Therefore, I urge the editor to ask the authors of the article to kindly extend their thoughts to the readers about how they came to their conclusions, to reveal the scientific sources for their statements, and to clarify contradicting conclusions. I think this is mandatory if we want to hold our journal to a higher standard than commercial magazines.

Carl Troilius
Malmö, Sweden

Reply to Troilius

It's a pleasure to reply to the letter of Dr. Troilius and have the opportunity to further clarify my paper.

Many papers have been published on the properties of glycolic acid, which is widely known to increase the thickness of both dermis and epidermis, enhancing the number of collagen fibers, of the dermis. As Cherie Ditre et al. [1] very extensively explained, daily applications of a lotion containing 25% alpha-hydroxy acids (glycolic, lactic or citric acid) causes an approximate 25% increase in skin thickness. The skin biopsies after 6 months showed that the epidermis was thicker, the papillary dermal changes included increased thickness, increased acid mucopolysaccharides, improved quality of the elastic fibers increased density of collagen. If Dr. Troilius wishes to go into this topic, I suggest that he read the following articles [1–7]. In my article I did not state that the collagen is produced in two to four weeks, but that as a rule we encourage glycolic acid use in all patients undergoing liposuction. I did not state that in two to four weeks the skin will be smoothened and the elastic fibers remodelled, but that the use of these products would stimulate the skin retraction capability through the stimulation of dermal collagen production and elastic fiber remodelling. I clearly explained in my paper that “this treatment is mandatory when the skin is hypotonic, and should start some months before the operation.” Therefore, Dr. Troilius did not read carefully the preoperative care paragraph in the section “Material and Method.”

About high protein diet and skin rejuvenation, I suggest review of the paper published by Dr. Romulo Mene that showed impressive results of skin retraction with the oral use of powdered protein extracted from selected young bovine dermis supplemented with amino acids and albumin [9].

With regard to the vitamin C therapy, it has been recommended by Zocchi [10] 1000 mg/day for three months after internal ultrasound liposuction. We believe that the external ultrasound carries “virtually no risk of free radical formation.” We used the word “Virtually” because we did not carry any study to prove this statement. So we do not really know if Vitamin C is indispensable or not but we still feel that the Zocchi-recommended dose of vitamin C may be prescribed with no added risk.

When general anesthesia is chosen, its level should be properly deep otherwise the blood rate and blood pressure may increase causing excessive bleeding. This is a basic principle of anesthesia. As excessive bleeding may cause uncontrolled consumption of coagulative factors, uncontrolled bleeding may occur.

We never stated that uncontrolled bleeding may occur in minimal liposuction.

Peter Bela Fodor, not I, named “superwet infiltration” the one that equals the amount of fat to be aspirated [11].

About the Lidocaine concentration, we do not use more then 20mg/Kg to avoid risk of Lidocaine toxicity. Thus, in local anesthesia we do small liposuctions where we presume to aspirate not more than 500 g of fat.

Regarding the ultrasound (US) conduction through gel, Dr. Troilius wonders if “any conduction of US can take place” through 2–3 cm of ultrasonic gel. The conduction of the US is enhanced by interposing gel between the probe and the skin. This is widely known and the ultrasonography techniques are full of practical examples of interposing a variety of materials like gel, epoxy resin, or paraffin. The thickness of 2–3 cm of gel does not impair the US transmission because the speed of US wave propagation in water (and gel is an aqueous medium) is very high (1410 m/sec). The use of gel is necessary to be sure no air is interposed between the probe and the skin, because the speed of US wave propagation in air is much slower (331 m/sec). Regarding why we used this thickness, there were practical reasons: with the US machine we had at the time when the article was written, the probe was not sterilizable, and we had to be sure that there was no air between the probe and the skin that would impair the US transmission. So we put the gel inside a sterile plastic cover retaining it in place with a rubber cover that would hold the gel in place. This rubber cover kept the gel very close to the probe with no air interposed, and the thickness of gel was approximately 2–3 cm.

We wrote that “during delivering external ultrasound the skin becomes slightly warmer,” which means a slight raise of the skin temperature, which is not relevant in regard of thermal damage.

The skin temperature rises because ultrasound is a source of energy. For all the reasons explained at point above, we never arrived at contradicting conclusions.

I trust that Dr. Troilius may be satisfied with the answers I gave.

Carlo Gasperoni
Rome, Italy

References

1. Ditre CM, Griffin TD, Murphy GF, Sueki H, Telegan B, Johnson WC, Yu RJ, Van Scott EJ: Effects of alpha-hydroxy acids on photoaged skin: a pilot clinical, histologic, and ultrastructural study. *J Am Acad Dermatol* **34**:187–195, 1996
2. Griffin TD, Murphy GF, Sueki H, Telegan B, Johnson WC, Ditre CM, Yu RJ, Van Scott EJ: Increased factor XIIIa transglutamine expression in dermal dendrocytes after treatment with alpha-hydroxyacids: potential physiologic significance. *J Am Acad Dermatol* **34**:196–203, 1996

3. Bernstein EF, Underhill CB, Lakkakorpi J, Ditre CM, Uitto J, Yu RJ, Scott EV: Citric acid increases viable epidermal thickness and glycosaminoglycan content of sun-damaged skin. *Dermatol Surg* **23**:689–694, 1997
4. Van Scott EJ, Ditre CM, Yu RJ: Alpha-hydroxyacids in the treatment of signs of photoaging. *Clin Dermatol*. **14**:217–226, 1996
5. Van Sgott EJ, Yu RJ: Alpha hydroxy acids: procedures for use in clinical practice. *Cutis* **43**:222–228, 1989
6. Newman N, Newman A, Moy LS, Babapour R, Harris AG, Moy RL: Clinical improvement of photoaged skin with 50% glycolic acid: a double-blind vehicle-controlled study. *Dermatol Surg* **22**:455–460, 1996
7. DiNardo JC, Grove GL, Moy LS: Clinical and histological effects of glycolic acid at different concentrations and pH levels. *Dermatol Surg* **22**:421–424, 1996
8. Moy LS, Howe K, Moy RL: Glycolic acid modulation of collagen production in human skin fibroblast cultures in vitro. *Dermatol Surg* **22**:439–441, 1996
9. Mene R: Rejuvenescimento: uso oral de colgeno em p (proteina isolada). *Abstracts, Int Symp on Recent Advances in Plastic Surgery*. S. Paulo, March 1989
10. Zocchi ML: Ultrasonic assisted lipoplasty. *Clin Plast Surg* **23**:575–598, 1996
11. Bela Fodor P: Defining wetting solutions in lipoplasty. *Plast Reconstr Surg* **103**:1519–1520, 1999

Coblation in Aesthetic Facial Rejuvenation

Recently, we had a chance to read Dr. Mancini's amazing article that was published in your journal

(Mancini, P.F. Coblation: a new technology and technique for skin resurfacing and other aesthetic surgical procedures, *Aesth Plast Surg* **25**(5):372–7 (2001). Since Cushing's great contribution to surgery in 1926, researchers constructed a variety of devices which used electricity as a means to heat tissue and control bleeding over the years. One of these constructions, *coblation* (which stands for "cold ablation") is a process employing bipolar radiofrequency technology to achieve tissue removal with minimal thermal damage to collateral tissue. Applications of this new technology in aesthetic facial surgery give promising results, like the ones in Dr. Mancini's paper. Dr. Mancini states that the device is adjusted in volts, which is the wrong parameter, in our opinion. Volt is the unit of measurement for voltage, the force that pushes electric current through impedance. The adjustable unit of the electrosurgical generators is the power, the amount of energy produced per second. And coblation device is actually a modified electrosurgical unit, as already mentioned by the author. We will be glad to learn if the voltage of the coblation is really adjustable. Otherwise, we think it would be proper to replace the "volts" by "watts," the unit of measurement for power.

Yener Demyrta
Reha Yavuzer