Erratum to: Preoperative Biliary Drainage of Severely Jaundiced Patients Increases Morbidity of Pancreaticoduodenectomy: Results of a Case-Control Study

Nikolaos Arkadopoulos · Maria A. Kyriazi ·
Ioannis S. Papanikolaou · Pantelis Vasiliou ·
Kassiani Theodoraki · Christos Lappas ·
Nikolaos Oikonomopoulos · Vassilios Smyrniotis

Published online: 8 November 2014
© Société Internationale de Chirurgie 2014

Erratum to: World J Surg (2014) 38:2967–2972
DOI 10.1007/s00268-014-2669-x

In the original article, percentages for post-pancreaticoduodenectomy morbidity and mortality are reversely cited

in the Introduction section, while the correct morbidity rates for the two groups are 24 versus 36 % (no PBD vs. PBD group), as reported in our text and not those erroneously provided in Table 2.

The online version of the original article can be found under doi:[10.1007/s00268-014-2669-x](https://doi.org/10.1007/s00268-014-2669-x).

N. Arkadopoulos · M. A. Kyriazi (✉) · P. Vasiliou · C. Lappas · V. Smyrniotis
4th Department of Surgery, University of Athens Medical School, Attikon Hospital, 1 Rimini Str, Chaidari, 12462 Athens, Greece
e-mail: mkyriazi@otenet.gr

I. S. Papanikolaou
Hepatogastroenterology Unit, 2nd Department of Internal Medicine, University of Athens Medical School, Attikon Hospital, Athens, Greece

K. Theodoraki
1st Department of Anaesthesiology, University of Athens Medical School, Aretaieion Hospital, Athens, Greece

N. Oikonomopoulos
2nd Department of Radiology, University of Athens Medical School, Attikon Hospital, Athens, Greece