ERRATUM

Erratum to: Safety and Hemostatic Effectiveness of the Fibrin Pad for Severe Soft-Tissue Bleeding During Abdominal, Retroperitoneal, Pelvic, and Thoracic (Non-cardiac) Surgery: A Randomized, Controlled, Superiority Trial

Jonathan Koea^{1,6} · Peter Baldwin² · Jessica Shen³ · B. Patel³ · Jonathan Batiller³ · Axel Arnaud³ · James Hart³ · Jeffrey Hammond³ · Craig Fischer⁴ · O. James Garden⁵

Published online: 28 July 2015
© Société Internationale de Chirurgie 2015

Erratum to: World J Surg DOI 10.1007/s00268-015-3106-5

The first sentence in the Results section of the Abstract is corrected as follows:

At 4 min post-randomization, 50/59 (84.7 %) subjects in the FP group and 10/32 (31.3 %) subjects in the SoC group

achieved hemostasis without needing re-treatment ($P < 0.0001$).

The correct first row entry in Table 3 is Hemostasis at 4 minutes without re-treatment, and the correct standard of treatment for this entry is 10/32 (31.3 %).

The online version of the original article can be found under doi:[10.1007/s00268-015-3106-5](https://doi.org/10.1007/s00268-015-3106-5).

✉ Jonathan Koea
jonathan.koea@waitematadhb.govt.nz

¹ Department of Surgery, Auckland City Hospital, Grafton, New Zealand

² Addenbrooke's Hospital, Cambridge, UK

³ Ethicon, Inc., Somerville, NJ, USA

⁴ Weill Cornell Medical College of Cornell University, Houston, TX, USA

⁵ Royal Infirmary of Edinburgh, Edinburgh, UK

⁶ Department of Surgery, North Shore Hospital, Private Bag 93505, Auckland 0620, New Zealand