Biologicals in rheumatology: Austrian experiences from a rheumatic outpatient clinic

C. Duftner · C. Dejaco · H. Larcher ·
M. Schirmer · M. Herold

Published online: 25 July 2008
© Springer-Verlag 2008

Erratum to: Rheumatol Int
DOI 10.1007/s00296-008-0621-9

Due to an unfortunate oversight Fig. 2 was rendered incorrectly. The correct Fig. 2 is given here:

Fig. 2 Disease duration (a) and age at onset of first biological therapy (b) remained constant over the observation period. *Whiskers* box plots show 50% of cases within the *boxes* and all data excluding mavericks between the end-points of the *whiskers* (*lines*). Differences were not significant according to the Kruskal–Wallis test

The online version of the original article can be found under
doi:[10.1007/s00296-008-0621-9](https://doi.org/10.1007/s00296-008-0621-9).

C. Duftner · C. Dejaco · H. Larcher · M. Schirmer · M. Herold
Clinical Department of Internal Medicine,
Division of General Internal Medicine,
Innsbruck Medical University, Innsbruck, Austria

C. Duftner (✉) · C. Dejaco · M. Schirmer
Department of Internal Medicine,
General Hospital of the Elisabethinen,
Voelkermärkerstrasse 15-19, 9020 Klagenfurt, Austria
e-mail: christina.duftner@ekh.at