Erratum to: Evaluation of the short-term efficacy of NSAIDs on patients with active ankylosing spondylitis in daily practice: a 3-month, longitudinal, observational study

Muhammet Cinar · Ayhan Dinc · Ismail Simsek · Hakan Erdem ·
Bayram Koc · Salih Pay · Selim Kilic

Published online: 19 March 2010
© Springer-Verlag 2010

Erratum to: Rheumatol Int (2010) 30:331–340 DOI 10.1007/s00296-009-0963-y

This article was unfortunately submitted with few errors in the issue 30(3) which we would like to correct here.

First; a sentence in the last paragraph of the results section on page 336: “Indometacin was given to 30.3% of patients, acemetacine was given to 28.6% of the patients, and other NSAIDs (diclofenac sodium, flurbiprofene, and celecoxib) were given to 26.7% of patients who responded to the treatment” causes misunderstanding.

The correct sentence is: “30.3, 28.6, and 26.7% of patients who treated with indometacin, acemetacin, and other NSAIDs (diclofenac sodium, flurbiprofene, celecoxib), in that order, had responded to their treatments.”

Second; the word “acemethazine” is incorrectly written at several lines in the text, however, the correct word should be as “acemetacin”.

The online version of the original article can be found under doi:[10.1007/s00296-009-0963-y](https://doi.org/10.1007/s00296-009-0963-y).

M. Cinar · B. Koc
Division of Internal Medicine,
Gulhane Military School of Medicine, Ankara, Turkey

A. Dinc · I. Simsek · H. Erdem · S. Pay
Division of Rheumatology,
Gulhane Military School of Medicine, Ankara, Turkey

S. Kilic
Department of Epidemiology and Public Health,
Gulhane Military School of Medicine, Ankara, Turkey

A. Dinc (✉)
GATA Romatoloji BD, Etlik, 06018 Ankara, Turkey
e-mail: adinc@gata.edu.tr; adinc@tr.net